arXiv:1103.2982v1 [math-ph] 15 Mar 2011

Addition theorems for spin spherical harmonics. I Preliminaries

Antonio O. Bouzas *
Departamento de Física Aplicada, CINVESTAV-IPN
Carretera Antigua a Progreso Km. 6, Apdo. Postal 73 "Cordemex"
Mérida 97310, Yucatán, México

January 14, 2013

Abstract

We develop a systematic approach to deriving addition theorems for, and some other bilocal sums of, spin spherical harmonics. In this first part we establish some necessary technical results. We discuss the factorization of orbital and spin degrees of freedom in certain products of Clebsch-Gordan coefficients, and obtain general explicit results for the matrix elements in configuration space of tensor products of arbitrary rank of the position and angular-momentum operators. These results are the basis of the addition theorems for spin spherical harmonics obtained in part II.

1 Introduction

The importance of the representation theory of the three-dimensional rotation group [1–7] in the study of all natural quantum systems hardly needs to be mentioned. That central role is due to the fact that the four fundamental interactions of nature, and their effective interactions relevant to nuclear, atomic and molecular physics, are all invariant under the rotation group. It is also for that reason that partial-wave expansions are essential tools in the analysis of both classical and quantum scattering processes.

The algebraic aspects of the computation of partial-wave expansions with spin states are best codified by the addition theorems for spin spherical harmonics. Spin-s spherical harmonics $Y^{\ell s}_{jj_z}(\widehat{r}\,)$ [4, 6] are the angular-momentum eigenfunctions relevant to the description of spin-s particles subject to spin-dependent central interactions.[1] As such, they are of interest in their own right beyond their above-mentioned specific application to partial-wave expansions that constitutes our main motivation for their study. In this paper, together with its second part [10] (hereafter referred to as II), we develop a systematic framework to derive addition theorems for spin spherical harmonics, and employ it to obtain those addition theorems for low spins, $0 < s', s \leq 3/2$, and for arbitrary integer spin s' if $s = 0$.

In this first part we establish the preliminary results that will be used as basic building blocks for constructing addition theorems for spin spherical harmonics in II. We consider first the factorization of orbital and spin degrees of freedom in products of Clebsch-Gordan (henceforth CG) coefficients of the form $\langle \ell', \ell'_z; s', s'_z | j, j_z\rangle\langle \ell, \ell_z; s, s_z | j, j_z\rangle$. We obtain a general factorization result, make its tensor and spinor structure explicit, and discuss the particular cases relevant to spins 1/2, 1 and 3/2. Such products of CG coefficients occur in the computation of physical quantities, such as scattering amplitudes or other matrix elements, related to transitions among states with definite orbital and spin quantum numbers. Thus, these results may be of interest independently of the addition theorems considered here and in II. We study also the matrix elements of arbitrary tensor products of orbital operators between angular-momentum projections of position eigenstates, for which we obtain completely general results. These are the other main ingredient in our derivation of addition theorems for spin spherical harmonics in II. Those matrix elements are closely related to bilocal spherical harmonics and other bilocal sums of ordinary spherical harmonics, as discussed in detail in II, so we expect that these results and the approach used to derive them should be applicable in other contexts as well.

*E-mail: abouzas@mda.cinvestav.mx

[1] Familiar examples of spin-s spherical harmonics are ordinary scalar spherical harmonics, $Y^{\ell 0}_{\ell\ell_z}(\widehat{r}\,) = Y_{\ell\ell_z}(\widehat{r}\,)$, and vector spherical harmonics [4, 6, 8, 9], $Y^{\ell 1}_{jj_z}(\widehat{r}\,)$.

The outline of the paper is as follows. In the next section we discuss the factorization of orbital and spin degrees of freedom in products of CG coefficients. In section 3 we introduce the angular-momentum projector operator and obtain general expressions for its matrix elements with orbital tensor operators. In appendix A we state our notation and conventions. In appendix B we give the definitions and main properties of the standard bases of irreducible tensors and spinors, of arbitrary spin, used as spin wavefunctions throughout the paper. A detailed list of general results for reduced matrix elements of arbitrary tensor products of orbital and spin operators, needed in the main sections of the paper, are given in appendix C. Finally, appendix D gathers some ancillary calculations needed in sect. 3.

2 Factorization of orbital and spin dependence

We begin by considering products of CG coefficients of the form

$$\begin{aligned} S(j,\ell',\ell,s',s;\ell'_z,\ell_z,s'_z,s_z) &= \sum_{j_z=-j}^{j} \langle \ell',\ell'_z;s',s'_z|j,j_z\rangle\langle \ell,\ell_z;s,s_z|j,j_z\rangle \\ &= \langle \ell',\ell'_z;s',s'_z|j,\ell'_z+s'_z\rangle\langle \ell,\ell_z;s,s_z|j,\ell_z+s_z\rangle\delta_{\ell'_z+s'_z,\ell_z+s_z} \ , \end{aligned} \tag{1}$$

with ℓ', ℓ integer and s', s integer or half-integer. We will usually write $S(j,\ell',\ell,s',s)$, omitting the last arguments for brevity. For the purpose of obtaining addition theorems for spin spherical harmonics we need to rewrite $S(j,\ell',\ell,s',s)$ as a sum of terms with a completely factorized dependence on orbital and spin angular-momentum projections, and to make explicit the tensor structure of those terms. The first goal is achieved by making use of relation (A.1) to write (see appendix A for our notation and conventions),

$$\begin{aligned} S(j,\ell',\ell,s',s) &= (-1)^{s'_z+s_z}(-1)^{\ell+s'-j}\frac{2j+1}{\sqrt{(2\ell'+1)(2s'+1)}}\sum_{\Delta=\Delta_{\min}}^{\Delta_{\max}}(2\Delta+1)\left\{\begin{array}{ccc}\ell' & \Delta & \ell\\ s & j & s'\end{array}\right\} \\ &\quad\times\langle \ell,\ell_z;\Delta,\Delta\ell_z|\ell',\ell'_z\rangle\langle s,s_z;\Delta,\Delta s_z|s',s'_z\rangle \ , \\ \Delta_{\min} &= \max\{|\Delta\ell|,|\Delta s|\} \ , \qquad \Delta_{\max} = \min\{\ell'+\ell,s'+s\} \ , \qquad \Delta\ell_z+\Delta s_z = 0 \ . \end{aligned} \tag{2}$$

In order to make the tensor structure of each term on the r.h.s. explicit, we use the Wigner-Eckart (henceforth WE) theorem [1–6, 8] to write the CG coefficients in terms of appropriate irreducible tensor operators,

$$\begin{aligned} S(j,\ell',\ell,s',s) &= \sum_{\Delta=\Delta_{\min}}^{\Delta_{\max}} C^{s's\Delta}_{\ell'\ell j}\langle \ell',\ell'_z|\widehat{\varepsilon}^{i_1\ldots i_\Delta}(\Delta\ell_z)\widehat{r}^{\,i_1}\ldots\widehat{r}^{\,i_{|\Delta\ell|}}L^{i_{|\Delta\ell|+1}}\ldots L^{i_\Delta}|\ell,\ell_z\rangle \\ &\quad\times\langle s',s'_z|\widehat{\varepsilon}^{h_1\ldots h_\Delta}(\Delta\ell_z)^*T^{h_1}\ldots T^{h_{|\Delta s|}}S^{h_{|\Delta s|+1}}\ldots S^{h_\Delta}|s,s_z\rangle \ , \\ C^{s's\Delta}_{\ell'\ell j} &= (-1)^{2s'_z}(-1)^{\ell+s'-j}\frac{2j+1}{\sqrt{(2\ell'+1)(2s'+1)}}(2\Delta+1)\left\{\begin{array}{ccc}\ell' & \Delta & \ell\\ s & j & s'\end{array}\right\} \\ &\quad\times\left(\langle\ell'||\widehat{\varepsilon}^{j_1\ldots j_\Delta}\widehat{r}^{\,j_1}\ldots\widehat{r}^{\,j_{|\Delta\ell|}}L^{j_{|\Delta\ell|+1}}\ldots L^{j_\Delta}||\ell\rangle\langle s'||\widehat{\varepsilon}^{k_1\ldots k_\Delta}T^{k_1}\ldots T^{k_{|\Delta s|}}S^{k_{|\Delta s|+1}}\ldots S^{k_\Delta}||s\rangle\right)^{-1} \ , \end{aligned} \tag{3}$$

where $\Delta_{\min,\max}$ are as in (2), and our notation for the standard basis tensors $\widehat{\varepsilon}^{i_1\ldots i_n}$ and for irreducible tensor operators is explained in appendix B. The operators appearing in (3) are the orbital $\vec{L}$ and spin $\vec{S}$ angular momenta, the position versor $\widehat{r}=\vec{r}/r$, and the spin-transition operator $\vec{T}$ defined here as the spin analog of $\widehat{r}$ (see sect. 2.4 below for a detailed discussion of this operator). The reduced matrix elements appearing in (3) are explicitly given for arbitrary values of their parameters in appendix C.

Whereas (3) is completely general, it is not yet explicit enough for the purpose of deriving addition theorems for spin spherical harmonics. From a practical point of view, we need only consider low values of s', s since quantum states with high values of spin occur infrequently in nature. In this paper we restrict ourselves to s', $s=0$, 1/2, 1, 3/2, and $|\Delta s|\leq 1$. For those spins the coefficients $C^{s's\Delta}_{\ell'\ell j}$ can be drastically simplified by exploiting the fact that $j-\ell'$ and $j-\ell$ can take only a small set of values. The spin matrix elements in (3) can be compactly evaluated in terms of standard tensors and spinors, and spin matrices. (The orbital matrix elements could, in principle, also be expressed in that way, but it would be impractical because ℓ', ℓ are not bounded above thus requiring tensor wave functions of arbitrarily large rank. We evaluate those matrix elements with a completely different technique below in section 3.) Clearly, we must

discuss separately the different values of s' and s, as those will determine the kind of spin spherical harmonics involved in the addition theorem. For fixed s' and s, different values of $\Delta\ell$ correspond to different addition theorems, so we must consider those cases separately as well. Furthermore, both in addition theorems for spin spherical harmonics and in partial wave expansions, terms with different $|\Delta\ell|$ have different tensor and spin structure.

2.1 $s' = 1/2 = s$

For $s' = 1/2 = s$ there are two possible values of $|\Delta\ell| = 0,\, 1$.

$\Delta\ell = 0$ In the case $s = 1/2$, j can take only two values and (3) reduces to

$$\begin{gathered} S(j,\ell,\ell,1/2,1/2) = C^{\frac12\frac12 0}_{\ell\ell j}\delta_{\ell'_z\ell_z}\delta_{s'_z s_z} + C^{\frac12\frac12 1}_{\ell\ell j}\langle \ell,\ell'_z|\widehat{\varepsilon}^k(\Delta\ell_z)L^k|\ell,\ell_z\rangle\,\widehat{\varepsilon}^h(\Delta\ell_z)^*\widehat{\chi}_A(s'_z)^*\frac12\sigma^h_{AB}\widehat{\chi}_B(s_z)\ , \\ C^{\frac12\frac12 0}_{\ell\ell j} = \frac{j+1/2}{2\ell+1}\ , \qquad C^{\frac12\frac12 1}_{\ell\ell j} = \frac{2(j-\ell)}{\ell+1/2}\ , \end{gathered} \tag{4}$$

which is a well-known result (see e.g., [11]). Notice that, due to the conservation of j_z, the product $\widehat{\varepsilon}^k(\Delta\ell_z)\widehat{\varepsilon}^h(\Delta\ell_z)^*$ in (4) can be replaced by δ^{kh}. We quote here also the simplified form of (2) in this case,

$$S(j,\ell,\ell,1/2,1/2) = C^{\frac12\frac12 0}_{\ell\ell j}\delta_{\ell'_z\ell_z}\delta_{s'_z s_z} + \frac{\sqrt{3}}{2}\sqrt{\ell(\ell+1)}C^{\frac12\frac12 1}_{\ell\ell j}\langle \ell,\ell_z;1,\Delta\ell_z|\ell,\ell'_z\rangle\langle 1/2,s'_z;1,\Delta\ell_z|1/2,s_z\rangle\ , \tag{5}$$

which results from (4) by applying the WE theorem.

$|\Delta\ell| = 1$ In this case $S(j,\ell',\ell,1/2,1/2)$ can be non-vanishing only if $\ell' = \ell+1$ and $j = \ell+1/2$, or if $\ell' = \ell-1$ and $j = \ell-1/2$. Equation (3) takes the simplified form,

$$S(j,\ell_1,\ell,1/2,1/2) = C^{\frac12\frac12 1}_{\ell_1\ell j}\langle \ell_1,\ell'_z|\widehat{\varepsilon}^k(\Delta\ell)\widehat{r}^k|\ell,\ell_z\rangle\,\widehat{\varepsilon}^h(\Delta\ell)^*\widehat{\chi}_A(s'_z)^*\frac12\sigma^h_{AB}\widehat{\chi}_B(s_z)\ , \quad C^{\frac12\frac12 1}_{\ell_1\ell j} = -2\ , \tag{6}$$

which is appropriate to derive an addition theorem for spin-1/2 spherical harmonics. As in the previous case, the product of standard versors can be substituted by δ^{kh}. Similarly, (2) reduces to

$$S(j,\ell_1,\ell,1/2,1/2) = -\Delta\ell\sqrt{\frac32}\sqrt{\frac{2\langle\ell\rangle+1}{2\ell_1+1}}\langle \ell,\ell_z;1,\Delta\ell_z|\ell_1,\ell'_z\rangle\langle 1/2,s'_z;1,\Delta\ell_z|1/2,s_z\rangle\ . \tag{7}$$

As a side remark, we notice that in this case $(2\langle\ell\rangle+1)/(2\ell_1+1) = (2j+1)/(2j+1+\Delta\ell)$, which provides an alternate form for (7).

2.2 $s' = 1 = s$

The general result (3) is greatly simplified in the case of $S(j,\ell',\ell,1,1)$ because, for fixed ℓ, $j-\ell$ can take only the three values ±1, 0. The spin matrix elements are given by

$$\begin{aligned} \langle 1,s'_z|\widehat{\varepsilon}^h(\Delta\ell_z)^*S^h|1,s_z\rangle &= -i\widehat{\varepsilon}^h(\Delta\ell_z)^*\left(\widehat{\varepsilon}(s'_z)^*\wedge\widehat{\varepsilon}(s_z)\right)^h\ , \\ \langle 1,s'_z|\widehat{\varepsilon}^{h_1h_1}(\Delta\ell_z)^*S^{h_1}S^{h_2}|1,s_z\rangle &= -\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^*\widehat{\varepsilon}^{h_1}(s'_z)^*\widehat{\varepsilon}^{h_2}(s_z)\ . \end{aligned} \tag{8}$$

There are three possible values for $\Delta\ell = 0,\, 1,\, 2$.

$|\Delta\ell| = 0$ In this case (3) can be written as

$$\begin{gathered} S(j,\ell,\ell,1,1) = C^{110}_{\ell\ell j}\delta_{\ell'_z\ell_z}\delta_{s'_z s_z} - C^{111}_{\ell\ell j}\langle \ell,\ell'_z|\widehat{\varepsilon}^k(\Delta\ell_z)L^k|\ell,\ell_z\rangle\,\widehat{\varepsilon}^h(\Delta\ell_z)^*i\left(\widehat{\varepsilon}(s'_z)^*\wedge\widehat{\varepsilon}(s_z)\right)^h \\ - C^{112}_{\ell\ell j}\langle \ell,\ell'_z|\widehat{\varepsilon}^{k_1k_2}(\Delta\ell_z)L^{k_1}L^{k_2}|\ell,\ell_z\rangle\,\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^*\widehat{\varepsilon}^{h_1}(s'_z)^*\widehat{\varepsilon}^{h_2}(s_z)\ , \\ C^{110}_{\ell\ell j} = \frac13\frac{2j+1}{2\ell+1}\ , \quad C^{111}_{\ell\ell j} = \frac{2}{D_{\ell j}}(2j+1)((j-\ell)(j+\ell+1)+1)\ , \quad C^{112}_{\ell\ell j} = \frac{4}{D_{\ell j}}(2j+1)\ , \\ D_{\ell j} = (-1)^{j-\ell-1}\left((j-\ell)^2+1\right)(j+\ell+2)(j+\ell+1)(j+\ell)\ . \end{gathered} \tag{9}$$

This form of (3) is needed in the derivation of addition theorems for vector spherical harmonics. Notice that in the term multiplied by $C^{111}_{\ell\ell j}$ we can replace $\widehat{\varepsilon}^k(\Delta\ell_z)\widehat{\varepsilon}^h(\Delta\ell_z)^*$ by δ^{kh}, and in the term multiplied by $C^{112}_{\ell\ell j}$ we can rewrite the matrix elements as

$$\langle \ell, \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) L^{k_1} L^{k_2} | \ell, \ell_z \rangle \, \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\varepsilon}^{h_1}(s'_z)^* \widehat{\varepsilon}^{h_2}(s_z) = \langle \ell, \ell'_z | \frac{1}{2} L^{\{p} L^{q\}_0} | \ell, \ell_z \rangle \, \widehat{\varepsilon}^{p}(s'_z)^* \widehat{\varepsilon}^{q}(s_z) \ . \tag{10}$$

The relation (2) in this case reduces to

$$\begin{aligned} S(j,\ell,1) = {} & C^{110}_{\ell\ell j} \delta_{\ell'_z \ell_z} \delta_{s'_z s_z} + \sqrt{2}\sqrt{\ell(\ell+1)} C^{111}_{\ell\ell j} \langle \ell, \ell_z; 1, \Delta\ell_z | \ell, \ell'_z \rangle \langle 1, s'_z; 1, \Delta\ell_z | 1, s_z \rangle \\ & + \frac{\sqrt{10}}{6} \sqrt{\ell(\ell+1)} \sqrt{(2\ell-1)(2\ell+3)} C^{112}_{\ell\ell j} \langle \ell, \ell_z; 2, \Delta\ell_z | \ell, \ell'_z \rangle \langle 1, s'_z; 2, \Delta\ell_z | 1, s_z \rangle \ , \end{aligned} \tag{11}$$

with $C^{11\Delta}_{\ell\ell j}$ as in (9).

$|\Delta\boldsymbol{\ell}| = 1$ In this case $S(j,\ell',\ell,1,1)$ can be non-vanishing only if $\ell' = \ell+1$ and $j = \ell+1$, ℓ, or if $\ell' = \ell-1$ and $j = \ell$, $\ell-1$. With that discrete set of values for $j-\ell$, the spin matrix elements (8), and the reduced matrix elements in appendix C, from (3) we obtain

$$\begin{aligned} S(j,\ell_1,\ell,1,1) = {} & -C^{111}_{\ell_1 \ell j} \langle \ell_1, \ell'_z | \widehat{\varepsilon}^{k}(\Delta\ell_z) \widehat{r}^{k} | \ell, \ell_z \rangle \, i\widehat{\varepsilon}^{h}(\Delta\ell_z)^* (\widehat{\varepsilon}(s'_z)^* \wedge \widehat{\varepsilon}(s_z))^h \\ & - C^{112}_{\ell_1 \ell j} \langle \ell_1, \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) \widehat{r}^{k_1} L^{k_2} | \ell, \ell_z \rangle \, \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\varepsilon}^{h_1}(s'_z)^* \widehat{\varepsilon}^{h_2}(s_z) \ , \\ C^{111}_{\ell_1 \ell j} = {} & -\frac{1}{2\langle\ell\rangle+1} \sqrt{2j+1} \sqrt{2j - \langle\ell\rangle + 1/2} \ , \quad C^{112}_{\ell_1 \ell j} = -4 \frac{(j - \langle\ell\rangle)}{2\langle\ell\rangle+1} \sqrt{\frac{2j+1}{2j - \langle\ell\rangle + 1/2}} \ . \end{aligned} \tag{12}$$

As before, on the first line of (12) we can replace $\widehat{\varepsilon}^{k}(\Delta\ell_z)\widehat{\varepsilon}^{h}(\Delta\ell_z)^*$ by δ^{kh}, and on the second line we can rewrite the matrix elements as

$$\langle \ell', \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) \widehat{r}^{k_1} L^{k_2} | \ell, \ell_z \rangle \, \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\varepsilon}^{h_1}(s'_z)^* \widehat{\varepsilon}^{h_2}(s_z) = \langle \ell', \ell'_z | \frac{1}{2} \widehat{r}^{\{p} L^{q\}_0} | \ell, \ell_z \rangle \, \widehat{\varepsilon}^{p}(s'_z)^* \widehat{\varepsilon}^{q}(s_z) \ . \tag{13}$$

The form of (2) in this case is

$$\begin{aligned} S(j,\ell_1,\ell,1,1) = {} & \Delta\ell \sqrt{\frac{2\langle\ell\rangle+1}{2\ell_1+1}} \Bigg(C^{111}_{\ell_1 \ell j} \langle \ell, \ell_z; 1, \Delta\ell_z | \ell_1, \ell'_z \rangle \langle 1, s'_z; 1, \Delta\ell_z | 1, s_z \rangle \\ & + \frac{1}{4} \sqrt{\frac{5}{3}} \sqrt{(2\langle\ell\rangle - 1)(2\langle\ell\rangle + 3)} C^{112}_{\ell_1 \ell j} \langle \ell, \ell_z; 2, \Delta\ell_z | \ell_1, \ell'_z \rangle \langle 1, s'_z; 2, \Delta\ell_z | 1, s_z \rangle \Bigg) \ , \end{aligned} \tag{14}$$

with the coefficients (12).

$|\Delta\boldsymbol{\ell}| = 2$ In this case $S(j,\ell',\ell,1,1)$ can be non-vanishing only if $\ell' = \ell+2$ and $j = \ell+1$, or if $\ell' = \ell-2$ and $j = \ell-1$. Using (8), (C.5) and (C.3), from (3) and (2) we obtain

$$\begin{aligned} S(j,\ell_2,\ell,1,1) = {} & -C^{112}_{\ell_2 \ell j} \langle \ell_2, \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) \widehat{r}^{k_1} \widehat{r}^{k_2} | \ell, \ell_z \rangle \, \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\varepsilon}^{h_1}(s'_z)^* \widehat{\varepsilon}^{h_2}(s_z) \\ = {} & \sqrt{\frac{5}{3}} \sqrt{\frac{j(j+1)}{(2j+1)(2\ell_2+1)}} C^{112}_{\ell_2 \ell j} \langle \ell, \ell_z; 2, \Delta\ell_z | \ell_2, \ell'_z \rangle \langle 1, s'_z; 2, \Delta\ell_z | 1, s_z \rangle \ , \\ C^{112}_{\ell_2 \ell j} = {} & \frac{2j+1}{\sqrt{j(j+1)}} \ . \end{aligned} \tag{15}$$

The first line is the form needed for an addition theorem for vector spherical harmonics. Notice that the matrix element on that line is traceless, because $\ell' \neq \ell$, and symmetric, so we can replace $\widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z)\widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^*$ there by $\delta^{k_1 h_1}\delta^{k_2 h_2}$. The second line is just (2).

2.3 $s = 3/2 = s'$

The spin matrix elements entering (3) when $s = 3/2 = s'$ are given by

$$\begin{aligned}
\widehat{\varepsilon}^{k}(\Delta\ell_z)^*\langle 3/2, s'_z|S^k|3/2, s_z\rangle &= \frac{3}{2}\widehat{\varepsilon}^{k}(\Delta\ell_z)^*\widehat{\chi}^i_A(s'_z)^*\sigma^k_{AB}\widehat{\chi}^i_B(s_z)\ ,\\
\widehat{\varepsilon}^{ij}(\Delta\ell_z)^*\langle 3/2, s'_z|S^iS^j|3/2, s_z\rangle &= -3\widehat{\varepsilon}^{ij}(\Delta\ell_z)^*\widehat{\chi}^i_A(s'_z)^*\widehat{\chi}^j_A(s_z)\ ,\\
\widehat{\varepsilon}^{ijk}(\Delta\ell_z)^*\langle 3/2, s'_z|S^iS^jS^k|3/2, s_z\rangle &= -\frac{3}{2}\widehat{\varepsilon}^{ijk}(\Delta\ell_z)^*\widehat{\chi}^i_A(s'_z)^*\sigma^j_{AB}\widehat{\chi}^k_B(s_z)\ .
\end{aligned} \tag{16}$$

The orbital angular momentum change is $0 \le \Delta\ell \le 3$.

$|\Delta\boldsymbol{\ell}| = \mathbf{0}$ Substituting (16) in (3) we get

$$\begin{aligned}
S(j,\ell,\ell,3/2,3/2) = {}& C^{\frac{3}{2}\frac{3}{2}0}_{\ell\ell j}\delta_{\ell'_z\ell_z}\delta_{s'_zs_z} + \frac{3}{2}C^{\frac{3}{2}\frac{3}{2}1}_{\ell\ell j}\langle\ell,\ell'_z|\widehat{\varepsilon}^{i}(\Delta\ell_z)L^i|\ell,\ell_z\rangle\widehat{\varepsilon}^{k}(\Delta\ell_z)^*\widehat{\chi}^h_A(s'_z)^*\sigma^k_{AB}\widehat{\chi}^h_B(s_z)\\
&- 3C^{\frac{3}{2}\frac{3}{2}2}_{\ell\ell j}\langle\ell,\ell'_z|\widehat{\varepsilon}^{i_1i_2}(\Delta\ell_z)L^{i_1}L^{i_2}|\ell,\ell_z\rangle\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\widehat{\chi}^{h_2}_A(s_z)\\
&- \frac{3}{2}C^{\frac{3}{2}\frac{3}{2}3}_{\ell\ell j}\langle\ell,\ell'_z|\widehat{\varepsilon}^{i_1i_2i_3}(\Delta\ell_z)L^{i_1}L^{i_2}L^{i_3}|\ell,\ell_z\rangle\widehat{\varepsilon}^{h_1h_2h_3}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\sigma^{h_2}_{AB}\widehat{\chi}^{h_3}_B(s_z)\ ,
\end{aligned} \tag{17a}$$

where the coefficients $C^{\frac{3}{2}\frac{3}{2}\Delta}_{\ell\ell j}$ given in (3) in this case take the form

$$\begin{aligned}
C^{\frac{3}{2}\frac{3}{2}0}_{\ell\ell j} &= \frac{1}{4}\,\frac{2j+1}{2\ell+1}\ ,\\
C^{\frac{3}{2}\frac{3}{2}1}_{\ell\ell j} &= \frac{2}{5}\,\frac{j-\ell}{2\ell+1}\left(j+\frac{1}{2}+4\left(j-\ell-\frac{3}{2}\right)(j-\ell)\left(j-\ell+\frac{3}{2}\right)\right)\frac{1}{\widetilde{D}_{\ell j}}\ ,\\
C^{\frac{3}{2}\frac{3}{2}2}_{\ell\ell j} &= -\frac{1}{2}\,\frac{1}{2\ell+1}\left(\frac{5}{4}-(j-\ell)^2\right)\left(j+\frac{1}{2}-2(j-\ell)^3+\frac{9}{2}(j-\ell)\right)\frac{1}{D_{\ell j}}\ ,\\
C^{\frac{3}{2}\frac{3}{2}3}_{\ell\ell j} &= -\frac{1}{2}\,\frac{(-1)^{j-\ell-1/2}}{|j-\ell|(2\ell+1)}\,\frac{1}{D_{\ell j}}\ ,\\
\widetilde{D}_{\ell j} &= \left(\ell+1+\frac{2}{3}\left(j-\ell-\frac{3}{2}\right)\left(j-\ell+\frac{1}{2}\right)(j-\ell+1)\right)\ ,\\
D_{\ell j} &= \widetilde{D}_{\ell j}\times\left(j-\frac{2}{3}\left(j-\ell-\frac{3}{2}\right)\left(j-\ell+\frac{1}{2}\right)(j-\ell+1)\right)\ .
\end{aligned} \tag{17b}$$

As above, we notice that the matrix elements in (17a) can also be written as,

$$\begin{aligned}
\widehat{\varepsilon}^{p}(\Delta\ell_z)\langle\ell,\ell'_z|L^p|\ell,\ell_z\rangle\widehat{\varepsilon}^{k}(\Delta\ell_z)^*\widehat{\chi}^i_A(s'_z)^*\sigma^k_{AB}\widehat{\chi}^i_B(s_z) &= \langle\ell,\ell'_z|L^k|\ell,\ell_z\rangle\widehat{\chi}^i_A(s'_z)^*\sigma^k_{AB}\widehat{\chi}^i_B(s_z)\ ,\\
2\widehat{\varepsilon}^{pq}(\Delta\ell_z)\langle\ell,\ell'_z|L^pL^q|\ell,\ell_z\rangle\widehat{\varepsilon}^{ij}(\Delta\ell_z)^*\widehat{\chi}^i_A(s'_z)^*\widehat{\chi}^j_B(s_z) &= \langle\ell,\ell'_z|L^{\{i}L^{j\}_0}|\ell,\ell_z\rangle\widehat{\chi}^i_A(s'_z)^*\widehat{\chi}^j_A(s_z)\ ,\\
6\widehat{\varepsilon}^{pqr}(\Delta\ell_z)\langle\ell,\ell'_z|L^pL^qL^r|\ell,\ell_z\rangle\widehat{\varepsilon}^{ijk}(\Delta\ell_z)^*\widehat{\chi}^i_A(s'_z)^*\sigma^k_{AB}\widehat{\chi}^j_B(s_z) &= \langle\ell,\ell'_z|L^{\{i}L^jL^{k\}_0}|\ell,\ell_z\rangle\\
&\quad\times\widehat{\chi}^i_A(s'_z)^*\sigma^j_{AB}\widehat{\chi}^k_B(s_z)\ .
\end{aligned} \tag{18}$$

With the matrix elements written as in (17a), $S(j,\ell,\ell,3/2,3/2)$ takes the form needed in the derivation of addition theorems for spin-3/2 spherical harmonics.

Similarly, (2) can be rewritten as

$$\begin{aligned}
S(j,\ell,\ell,3/2,3/2) = {}& \kappa^{\frac{3}{2}\frac{3}{2}0}_{\ell\ell j}\delta_{\ell'_z\ell_z}\delta_{s'_zs_z} + \kappa^{\frac{3}{2}\frac{3}{2}1}_{\ell\ell j}\langle\ell,\ell_z;1,\Delta\ell_z|\ell,\ell'_z\rangle\langle 3/2,s'_z;1,\Delta\ell_z|3/2,s_z\rangle\\
&+ \kappa^{\frac{3}{2}\frac{3}{2}2}_{\ell\ell j}\langle\ell,\ell_z;2,\Delta\ell_z|\ell,\ell'_z\rangle\langle 3/2,s'_z;2,\Delta\ell_z|3/2,s_z\rangle\\
&+ \kappa^{\frac{3}{2}\frac{3}{2}3}_{\ell\ell j}\langle\ell,\ell_z;3,\Delta\ell_z|\ell,\ell'_z\rangle\langle 3/2,s'_z;3,\Delta\ell_z|3/2,s_z\rangle\ ,\\
\kappa^{\frac{3}{2}\frac{3}{2}0}_{\ell\ell j} = C^{\frac{3}{2}\frac{3}{2}0}_{\ell\ell j},\quad \kappa^{\frac{3}{2}\frac{3}{2}1}_{\ell\ell j} &= \sqrt{\frac{15}{4}}\sqrt{\ell(\ell+1)}C^{\frac{3}{2}\frac{3}{2}1}_{\ell\ell j},\quad \kappa^{\frac{3}{2}\frac{3}{2}2}_{\ell\ell j} = \sqrt{\frac{5}{4}}\sqrt{\ell(\ell+1)}\sqrt{(2\ell-1)(2\ell+3)}C^{\frac{3}{2}\frac{3}{2}2}_{\ell\ell j},\\
\kappa^{\frac{3}{2}\frac{3}{2}3}_{\ell\ell j} &= \frac{3}{2}\sqrt{\frac{7}{20}}\sqrt{(\ell-1)\ell(\ell+1)(\ell+2)}\sqrt{(2\ell-1)(2\ell+3)}C^{\frac{3}{2}\frac{3}{2}3}_{\ell\ell j}\ ,
\end{aligned} \tag{19}$$

Eqs. (17) and (19) are of course related by the WE theorem, with the reduced matrix elements in appendix C.

$|\Delta\boldsymbol{\ell}| = \mathbf{1}$ In this case $S(j,\ell',\ell,3/2,3/2)$ can be non-vanishing only if $\ell' = \ell+1$ and $j = \ell+3/2$, $\ell+1/2$, $\ell-1/2$, or if $\ell' = \ell-1$ and $j = \ell+1/2$, $\ell-1/2$, $\ell-3/2$. Evaluating the spin matrix elements with (16) we find,

$$\begin{aligned} S(j,\ell_1,\ell,3/2,3/2) = {} & \frac{3}{2} C^{\frac{3}{2}\frac{3}{2}1}_{\ell_1\ell j} \langle \ell_1,\ell'_z|\widehat{\varepsilon}^k(\Delta\ell_z)\widehat{r}^k|\ell,\ell_z\rangle\, \widehat{\varepsilon}^h(\Delta\ell_z)^* \widehat{\chi}^j_A(s'_z)^* \sigma^h_{AB} \widehat{\chi}^j_B(s_z) \\ & - 3 C^{\frac{3}{2}\frac{3}{2}2}_{\ell_1\ell j} \langle \ell_1,\ell'_z|\widehat{\varepsilon}^{i_1 i_2}(\Delta\ell_z)\widehat{r}^{i_1} L^{i_2}|\ell,\ell_z\rangle\, \widehat{\varepsilon}^{j_1 j_2}(\Delta\ell_z)^* \widehat{\chi}^{j_1}_A(s'_z)^* \widehat{\chi}^{j_2}_A(s_z) \\ & - \frac{3}{2} C^{\frac{3}{2}\frac{3}{2}3}_{\ell_1\ell j} \langle \ell_1,\ell'_z|\widehat{\varepsilon}^{i_1 i_2 i_3}(\Delta\ell_z)\widehat{r}^{i_1} L^{i_2} L^{i_3}|\ell,\ell_z\rangle\, \widehat{\varepsilon}^{j_1 j_2 j_3}(\Delta\ell_z)^* \widehat{\chi}^{j_1}_A(s'_z)^* \sigma^{j_2}_{AB} \widehat{\chi}^{j_3}_B(s_z)\ , \end{aligned} \tag{20a}$$

with the coefficients $C^{\frac{3}{2}\frac{3}{2}\Delta}_{\ell_1\ell j}$ given by

$$\begin{aligned} C^{\frac{3}{2}\frac{3}{2}1}_{\ell_1\ell j} &= -\frac{1}{5}\,\frac{j+1/2}{2\langle\ell\rangle+1}\,\frac{\sqrt{\left(j+\langle\ell\rangle+1\right)^2-4}\,\sqrt{4-\left(j-\langle\ell\rangle\right)^2}}{\sqrt{\langle\ell\rangle(\langle\ell\rangle+1)}}\ , \\ C^{\frac{3}{2}\frac{3}{2}2}_{\ell_1\ell j} &= -2\frac{\Big(3\langle\ell\rangle-j+1\Big)}{(2\langle\ell\rangle+1)\sqrt{\langle\ell\rangle(\langle\ell\rangle+1)}}\,\frac{\Big((j-\langle\ell\rangle)(j+1/2)+1/2\Big)\Big(5(j-\langle\ell\rangle)^2-4\Big)}{\sqrt{(j+\langle\ell\rangle+1)^2-4}\sqrt{4-(j-\langle\ell\rangle)^2}}\ , \\ C^{\frac{3}{2}\frac{3}{2}3}_{\ell_1\ell j} &= -4\frac{\sqrt{\langle\ell\rangle(\langle\ell\rangle+1)}}{(2\langle\ell\rangle+1)j(j+1)}\,\frac{\Big(3j-\langle\ell\rangle+1\Big)\Big(3/2(j-\langle\ell\rangle)^2-1\Big)}{\sqrt{(j+\langle\ell\rangle+1)^2-4}\sqrt{4-(j-\langle\ell\rangle)^2}}\ . \end{aligned} \tag{20b}$$

This expression in terms of matrix elements of irreducible tensor operators is needed in the derivation of addition theorems for spin-3/2 spherical harmonics. We notice also that the matrix elements in (20a) can be written without standard tensors as,

$$\begin{aligned} \langle \ell_1,\ell'_z|\widehat{\varepsilon}^k(\Delta\ell_z)\widehat{r}^k|\ell,\ell_z\rangle\, \widehat{\varepsilon}^h(\Delta\ell_z)^* \widehat{\chi}^j_A(s'_z)^* \sigma^h_{AB} \widehat{\chi}^j_B(s_z) &= \langle \ell_1,\ell'_z|\widehat{r}^k|\ell,\ell_z\rangle\, \widehat{\chi}^j_A(s'_z)^* \sigma^k_{AB} \widehat{\chi}^j_B(s_z)\ , \\ \langle \ell_1,\ell'_z|\widehat{\varepsilon}^{i_1 i_2}(\Delta\ell_z)\widehat{r}^{i_1} L^{i_2}|\ell,\ell_z\rangle\, \widehat{\varepsilon}^{j_1 j_2}(\Delta\ell_z)^* \widehat{\chi}^{j_1}_A(s'_z)^* \widehat{\chi}^{j_2}_A(s_z) &= \frac{1}{2}\langle \ell_1,\ell'_z|\widehat{r}^{\{i_1} L^{i_2\}}|\ell,\ell_z\rangle\, \widehat{\chi}^{i_1}_A(s'_z)^* \widehat{\chi}^{i_2}_A(s_z)\ , \\ \langle \ell_1,\ell'_z|\widehat{\varepsilon}^{i_1 i_2 i_3}(\Delta\ell_z)\widehat{r}^{i_1} L^{i_2} L^{i_3}|\ell,\ell_z\rangle\, \widehat{\varepsilon}^{j_1 j_2 j_3}(\Delta\ell_z)^* \widehat{\chi}^{j_1}_A(s'_z)^* \sigma^{j_2}_{AB} \widehat{\chi}^{j_3}_B(s_z) &= \frac{1}{6}\langle \ell_1,\ell'_z|\widehat{r}^{\{i_1} L^{i_2} L^{i_3\}_0}|\ell,\ell_z\rangle \\ &\quad \times \widehat{\chi}^{i_1}_A(s'_z)^* \sigma^{i_2}_{AB} \widehat{\chi}^{i_3}_B(s_z)\ . \end{aligned} \tag{21}$$

Analogously, (2) can be written as

$$\begin{aligned} S(j,\ell_1,\ell,3/2,3/2) = {} & \kappa^{\frac{3}{2}\frac{3}{2}1}_{\ell_1\ell j} \langle \ell,\ell_z;1,\Delta\ell_z|\ell_1,\ell'_z\rangle\langle 3/2,s'_z;1,\Delta\ell_z|3/2,s_z\rangle \\ & + \kappa^{\frac{3}{2}\frac{3}{2}2}_{\ell_1\ell j} \langle \ell,\ell_z;2,\Delta\ell_z|\ell_1,\ell'_z\rangle\langle 3/2,s'_z;2,\Delta\ell_z|3/2,s_z\rangle \\ & + \kappa^{\frac{3}{2}\frac{3}{2}3}_{\ell_1\ell j} \langle \ell,\ell_z;3,\Delta\ell_z|\ell_1,\ell'_z\rangle\langle 3/2,s'_z;3,\Delta\ell_z|3/2,s_z\rangle\ , \end{aligned} \tag{22a}$$

with,

$$\begin{aligned} \kappa^{\frac{3}{2}\frac{3}{2}1}_{\ell_1\ell j} &= \frac{1}{2}\sqrt{\frac{15}{2}}\,\frac{\Delta\ell}{\sqrt{2\ell_1+1}}\,\sqrt{2\langle\ell\rangle+1}\,C^{\frac{3}{2}\frac{3}{2}1}_{\ell_1\ell j}\ , \\ \kappa^{\frac{3}{2}\frac{3}{2}2}_{\ell_1\ell j} &= \frac{1}{4}\sqrt{\frac{15}{2}}\,\frac{\Delta\ell}{\sqrt{2\ell_1+1}}\,\sqrt{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)}\,C^{\frac{3}{2}\frac{3}{2}2}_{\ell_1\ell j}\ , \\ \kappa^{\frac{3}{2}\frac{3}{2}3}_{\ell_1\ell j} &= \frac{1}{8}\sqrt{\frac{21}{5}}\,\frac{\Delta\ell}{\sqrt{2\ell_1+1}}\,\sqrt{(2\langle\ell\rangle-2)(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)(2\langle\ell\rangle+4)}\,C^{\frac{3}{2}\frac{3}{2}3}_{\ell_1\ell j}\ , \end{aligned} \tag{22b}$$

where the coefficients $C^{\frac{3}{2}\frac{3}{2}\Delta}_{\ell_1\ell j}$ are given in (20b).

$|\Delta\boldsymbol{\ell}| = \mathbf{2}$ In this case $S(j,\ell',\ell,3/2)$ can be non-vanishing only if $\ell' = \ell+2$ and $j = \ell+3/2, \ell+1/2$ or $\ell' = \ell-2$ and $j = \ell-1/2, \ell-3/2$. Taking into account this reduced set of j values, and the spin matrix elements (16), eq. (3) takes the form

$$\begin{aligned} S(j,\ell_2,\ell,3/2,3/2) = &-3C^{\frac{3}{2}\frac{3}{2}2}_{\ell_2\ell j}\langle \ell_2,\ell'_z|\widehat{\varepsilon}^{k_1k_2}(\Delta\ell_z)\widehat{r}^{k_1}\widehat{r}^{k_2}|\ell,\ell_z\rangle\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\widehat{\chi}^{h_2}_A(s_z) \\ &-\frac{3}{2}C^{\frac{3}{2}\frac{3}{2}3}_{\ell_2\ell j}\langle \ell_2,\ell'_z|\widehat{\varepsilon}^{k_1k_2k_3}(\Delta\ell_z)\widehat{r}^{k_1}\widehat{r}^{k_2}L^{k_3}|\ell,\ell_z\rangle\widehat{\varepsilon}^{h_1h_2h_3}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\sigma^{h_2}_{AB}\widehat{\chi}^{h_3}_B(s_z)\ , \\ C^{\frac{3}{2}\frac{3}{2}2}_{\ell_2\ell j} = &\frac{2}{\sqrt{3}}\frac{3j-2\langle\ell\rangle+1/2}{\sqrt{(2j-1)(2j+3)}}\ , \quad C^{\frac{3}{2}\frac{3}{2}3}_{\ell_2\ell j} = \frac{8}{\sqrt{3}}\frac{j-\langle\ell\rangle}{\sqrt{(2j-1)(2j+3)}}\ . \end{aligned} \tag{23}$$

We note here that the matrix elements appearing in this equation can also be written as

$$\begin{aligned} &\langle \ell',\ell'_z|\widehat{\varepsilon}^{k_1k_2}(\Delta\ell_z)\widehat{r}^{k_1}\widehat{r}^{k_2}|\ell,\ell_z\rangle\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\widehat{\chi}^{h_2}_A(s_z) = \langle \ell',\ell'_z|\widehat{r}^{i_1}\widehat{r}^{i_2}|\ell,\ell_z\rangle\widehat{\chi}^{i_1}_A(s'_z)^*\widehat{\chi}^{i_2}_A(s_z)\ , \\ &\langle \ell',\ell'_z|\widehat{\varepsilon}^{k_1k_2k_3}(\Delta\ell_z)\widehat{r}^{k_1}\widehat{r}^{k_2}L^{k_3}|\ell,\ell_z\rangle\widehat{\varepsilon}^{h_1h_2h_3}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\sigma^{h_2}_{AB}\widehat{\chi}^{h_3}_B(s_z) = \\ &\qquad\qquad \frac{1}{6}\langle \ell',\ell'_z|\widehat{r}^{\{i_1}\widehat{r}^{i_2}L^{i_3\}_0}|\ell,\ell_z\rangle\widehat{\chi}^{i_1}_A(s'_z)^*\sigma^{i_2}_{AB}\widehat{\chi}^{i_3}_B(s_z)\ , \end{aligned} \tag{24}$$

where the matrix element in the r.h.s. of the first equality is traceless because $\ell' \neq \ell$. Similarly, from eq. (2) we get

$$\begin{aligned} S(j,\ell_2,\ell,3/2,3/2) = &\ \kappa^{\frac{3}{2}\frac{3}{2}2}_{\ell_2\ell j}\langle \ell,\ell_z;2,\Delta\ell_z|\ell_2,\ell'_z\rangle\langle 3/2,s'_z;2,\Delta\ell_z|3/2,s_z\rangle \\ &+\kappa^{\frac{3}{2}\frac{3}{2}3}_{\ell_2\ell j}\langle \ell,\ell_z;3,\Delta\ell_z|\ell_2,\ell'_z\rangle\langle 3/2,s'_z;3,\Delta\ell_z|3/2,s_z\rangle\ , \\ \kappa^{\frac{3}{2}\frac{3}{2}2}_{\ell_2\ell j} = &\sqrt{\frac{15}{2}}\sqrt{2\ell+1}\sqrt{\frac{\langle\ell\rangle(\langle\ell\rangle+1)}{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)}}C^{\frac{3}{2}\frac{3}{2}2}_{\ell_2\ell j}\ , \\ \kappa^{\frac{3}{2}\frac{3}{2}3}_{\ell_2\ell j} = &\frac{1}{2}\sqrt{\frac{21}{2}}\sqrt{2\ell+1}\sqrt{\frac{(\langle\ell\rangle-1)\langle\ell\rangle(\langle\ell\rangle+1)(\langle\ell\rangle+2)}{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)}}C^{\frac{3}{2}\frac{3}{2}3}_{\ell_2\ell j}\ , \end{aligned} \tag{25}$$

with the coefficients $C^{\frac{3}{2}\frac{3}{2}\Delta}_{\ell_2\ell j}$ of (23).

$|\Delta\boldsymbol{\ell}| = \mathbf{3}$ In this case $S(j,\ell',\ell,3/2,3/2)$ can be non-vanishing only if $\ell' = \ell+3$ and $j = \ell+3/2$, or $\ell' = \ell-3$ and $j = \ell-3/2$. Eq. (3) then reduces to

$$\begin{aligned} S(j,\ell_3,\ell,3/2,3/2) = &-\frac{3}{2}C^{\frac{3}{2}\frac{3}{2}3}_{\ell_3\ell j}\langle \ell_3,\ell'_z|\widehat{\varepsilon}^{k_1k_2k_3}(\Delta\ell_z)\widehat{r}^{k_1}\widehat{r}^{k_2}\widehat{r}^{k_3}|\ell,\ell_z\rangle \\ &\times\widehat{\varepsilon}^{h_1h_2h_3}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\sigma^{h_2}_{AB}\widehat{\chi}^{h_3}_B(s_z)\ , \\ C^{\frac{3}{2}\frac{3}{2}3}_{\ell_3\ell j} = &-\frac{8}{3}\sqrt{\frac{j(j+1)}{(2j-1)(2j+3)}}\ , \end{aligned} \tag{26}$$

which is the form needed to formulate an addition theorem for spin-3/2 spherical harmonics with $\ell' = \ell\pm 3$. The matrix element in (26) can also be written

$$\begin{aligned} &\langle \ell',\ell'_z|\widehat{\varepsilon}^{k_1k_2k_3}(\Delta\ell_z)\widehat{r}^{k_1}\widehat{r}^{k_2}\widehat{r}^{k_3}|\ell,\ell_z\rangle\widehat{\varepsilon}^{h_1h_2h_3}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\sigma^{h_2}_{AB}\widehat{\chi}^{h_3}_B(s_z) = \\ &\qquad\qquad = \frac{1}{6}\langle \ell',\ell'_z|\widehat{r}^{\{k_1}\widehat{r}^{k_2}\widehat{r}^{k_3\}_0}|\ell,\ell_z\rangle\widehat{\chi}^{k_1}_A(s'_z)^*\sigma^{k_2}_{AB}\widehat{\chi}^{k_3}_B(s_z)\ . \end{aligned} \tag{27}$$

Similarly, (2) reduces to

$$\begin{aligned} S(j,\ell_3,\ell,3/2,3/2) = &\ \kappa^{\frac{3}{2}\frac{3}{2}3}_{\ell_3\ell j}\langle \ell,\ell_z;3,\Delta\ell_z|\ell_3,\ell'_z\rangle\langle 3/2,s'_z;3,\Delta\ell_z|3/2,s_z\rangle\ , \\ \kappa^{\frac{3}{2}\frac{3}{2}3}_{\ell_3\ell j} = &\frac{\sqrt{7}}{32}\Delta\ell\sqrt{\frac{2\ell+1}{\langle\ell\rangle+2}}\sqrt{\frac{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)}{(\langle\ell\rangle-1)\langle\ell\rangle(\langle\ell\rangle+1)}}C^{\frac{3}{2}\frac{3}{2}3}_{\ell_3\ell j}\ , \end{aligned} \tag{28}$$

with $C^{\frac{3}{2}\frac{3}{2}3}_{\ell_3\ell j}$ from (26).

2.4 The spin transition operator

We define the spin transition operator $\vec{T}$ as a spin vector operator, commuting with all orbital operators, and satisfying

$$[S^i, T^j] = i\varepsilon^{ijk}T^k \ , \qquad \langle s'||\widehat{\varepsilon}^i T^i||s\rangle = \frac{s'-s}{\sqrt{2}}\sqrt{\frac{2\langle s\rangle+1}{2s'+1}}\left(\delta_{s'(s+1)}+\delta_{s'(s-1)}\right). \tag{29}$$

From (29) we have,

$$\langle s', s'_z|\vec{T}^2|s, s_z\rangle = \begin{cases} \delta_{s's}\delta_{s'_z s_z} & \text{if } s' > 1/2 \text{ or } s > 1/2 \\ \frac{3}{4}\delta_{s'_z s_z} & \text{if } s' = s = 1/2 \end{cases} . \tag{30}$$

From (29) and (30) the operator $\vec{T}$ is seen to be the spin-space analog of the orbital operator $\widehat{r}$. In particular, the reduced matrix elements of tensor products of $\widehat{r}$ and $\vec{L}$ given in Appendix C apply without changes to tensor products of $\vec{T}$ and $\vec{S}$.

2.5 $s', s = 1, 0$ or $0, 1$

The relevant spin matrix element in this case is,

$$\langle 1, s'_z|T^h|0, 0\rangle = \frac{1}{\sqrt{3}}\widehat{\varepsilon}^h(s'_z)^* \ . \tag{31}$$

$|\Delta\ell|$ can take the values 0 and 1.

$|\Delta\boldsymbol{\ell}| = 0$ $S(j, \ell, \ell, 1, 0)$ (resp. $S(j, \ell, \ell, 0, 1)$) can be non-vanishing only if $j = \ell$ (resp. $j = \ell'$). Then, (3) and (2) take the form,

$$\begin{gathered} S(\ell, \ell, \ell, 1, 0) = \frac{1}{\sqrt{3}}C^{101}_{\ell\ell\ell}\langle \ell, \ell'_z|L^k|\ell, \ell_z\rangle\widehat{\varepsilon}^k(s'_z)^* = -\sqrt{3}\langle \ell, \ell_z; 1, \Delta\ell_z|\ell, \ell'_z\rangle\langle 1, s'_z; 1, \Delta\ell_z|0, 0\rangle \ , \\ C^{101}_{\ell\ell\ell} = \sqrt{\frac{3}{\ell(\ell+1)}} \ . \end{gathered} \tag{32}$$

Similarly,

$$S(\ell, \ell, \ell, 0, 1) = \frac{1}{\sqrt{3}}C^{011}_{\ell\ell\ell}\langle \ell, \ell'_z|L^k|\ell, \ell_z\rangle\widehat{\varepsilon}^k(s_z) = \langle \ell, \ell_z; 1, \Delta\ell_z|\ell, \ell'_z\rangle\langle 0, 0; 1, \Delta\ell_z|1, s_z\rangle \ , \tag{33}$$

with $C^{011}_{\ell\ell\ell} = C^{101}_{\ell\ell\ell}$.

$|\Delta\boldsymbol{\ell}| = 1$ In this case $S(j, \ell', \ell, s', s)$ can be non-vanishing only if $j = \ell$ (if $s = 0$) or ℓ' (if $s' = 0$). Thus, with the spin matrix element (31) we get, from (3) and (2) ,

$$\begin{gathered} S(\ell, \ell_1, \ell, 1, 0) = \frac{C^{101}_{\ell_1\ell\ell}}{\sqrt{3}}\langle \ell_1, \ell'_z|\widehat{r}^k|\ell, \ell_z\rangle\widehat{\varepsilon}^k(s'_z)^* = \sqrt{3}\sqrt{\frac{2\ell+1}{2\ell_1+1}}\langle \ell, \ell_z; 1, \Delta\ell_z|\ell_1, \ell'_z\rangle\langle 1, s'_z; 1, \Delta\ell_z|0, 0\rangle \ , \\ C^{101}_{\ell_1\ell\ell} = -\sqrt{6}\Delta\ell\sqrt{\frac{2\ell+1}{2\langle\ell\rangle+1}} \ . \end{gathered} \tag{34}$$

Analogously,

$$\begin{gathered} S(\ell_1, \ell_1, \ell, 0, 1) = \frac{C^{011}_{\ell_1\ell\ell_1}}{\sqrt{3}}\langle \ell_1, \ell'_z|\widehat{r}^k|\ell, \ell_z\rangle\widehat{\varepsilon}^k(s'_z) = \langle \ell, \ell_z; 1, \Delta\ell_z|\ell_1, \ell'_z\rangle\langle 1, s'_z; 1, \Delta\ell_z|0, 0\rangle \ , \\ C^{011}_{\ell_1\ell\ell'} = \sqrt{6}\Delta\ell\sqrt{\frac{2\ell'+1}{2\langle\ell\rangle+1}} \ . \end{gathered} \tag{35}$$

2.6 $s', s = 3/2, 1/2$ or $1/2, 3/2$

The spin matrix elements appearing in (3) in this case are

$$\begin{aligned}\langle 3/2, s'_z|\widehat{\varepsilon}^h(\Delta\ell_z)^* T^h|1/2, s_z\rangle &= \frac{1}{2}\sqrt{\frac{3}{2}}\widehat{\varepsilon}^h(\Delta\ell_z)^*\widehat{\chi}^h_A(s'_z)^*\widehat{\chi}_A(s_z)\ ,\\ \langle 3/2, s'_z|\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^* T^{h_1}S^{h_2}|1/2, s_z\rangle &= \frac{1}{4}\sqrt{\frac{3}{2}}\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\sigma^{h_2}_{AB}\widehat{\chi}_B(s_z)\ .\end{aligned} \tag{36}$$

There are three possible values for $0 \le |\Delta\ell| \le 2$.

$|\Delta\boldsymbol{\ell}| = \mathbf{0}$ If $s' = 3/2$, $s = 1/2$, and $\ell' = \ell$, then $S(j, \ell, \ell, 3/2, 1/2)$ can be non-vanishing only if $j = \ell \pm 1/2$. In the case $s' = 1/2$, $s = 3/2$, it must be $j = \ell' \pm 1/2$. Substituting (36) in (3) we obtain

$$\begin{aligned}S(j, \ell, \ell, 3/2, 1/2) = &\frac{1}{2}\sqrt{\frac{3}{2}}C^{\frac{3}{2}\frac{1}{2}1}_{\ell\ell j}\langle \ell, \ell'_z|\widehat{\varepsilon}^k(\Delta\ell_z)L^k|\ell, \ell_z\rangle\widehat{\varepsilon}^h(\Delta\ell_z)^*\widehat{\chi}^h_A(s'_z)^*\widehat{\chi}_A(s_z)\\ &+\frac{1}{4}\sqrt{\frac{3}{2}}C^{\frac{3}{2}\frac{1}{2}2}_{\ell\ell j}\langle \ell, \ell'_z|\widehat{\varepsilon}^{k_1k_2}(\Delta\ell_z)L^{k_1}L^{k_2}|\ell, \ell_z\rangle\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\sigma^{h_2}_{AB}\widehat{\chi}_B(s_z)\ ,\end{aligned} \tag{37a}$$

with

$$C^{\frac{3}{2}\frac{1}{2}1}_{\ell\ell j} = \frac{2}{2\ell+1}\sqrt{\frac{2j-\ell+1/2}{2\ell-j+1/2}}\ , \qquad C^{\frac{3}{2}\frac{1}{2}2}_{\ell\ell j} = 8\frac{j-\ell}{2\ell+1}\frac{1}{\sqrt{(2j-\ell+1/2)(2\ell-j+1/2)}}\ . \tag{37b}$$

The matrix elements in (37a) can also be written without standard tensors as,

$$\begin{aligned}\langle \ell, \ell'_z|\widehat{\varepsilon}^k(\Delta\ell_z)L^k|\ell, \ell_z\rangle\widehat{\varepsilon}^h(\Delta\ell_z)^*\widehat{\chi}^h_A(s'_z)^*\widehat{\chi}_A(s_z) &= \langle \ell, \ell'_z|L^k|\ell, \ell_z\rangle\widehat{\chi}^k_A(s'_z)^*\widehat{\chi}_A(s_z)\ ,\\ \langle \ell, \ell'_z|\widehat{\varepsilon}^{k_1k_2}(\Delta\ell_z)L^{k_1}L^{k_2}|\ell, \ell_z\rangle\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^*\widehat{\chi}^{h_1}_A(s'_z)^*\sigma^{h_2}_{AB}\widehat{\chi}_B(s_z) &=\\ = \frac{1}{2}\langle \ell, \ell'_z|L^{\{k_1}L^{k_2\}_0}|\ell, \ell_z\rangle\widehat{\chi}^{k_1}_A(s'_z)^*\sigma^{k_2}_{AB}\widehat{\chi}_B(s_z)\ .\end{aligned} \tag{38}$$

Similarly, (2) takes the simpler form,

$$\begin{aligned}S(j, \ell, \ell, 3/2, 1/2) = &-\frac{\sqrt{3}}{2}\sqrt{\ell(\ell+1)}C^{\frac{3}{2}\frac{1}{2}1}_{\ell\ell j}\langle \ell, \ell_z; 1, \Delta\ell_z|\ell, \ell'_z\rangle\langle 3/2, s'_z; 1, \Delta\ell_z|1/2, s_z\rangle\\ &-\frac{\sqrt{5}}{8}\sqrt{\ell(\ell+1)}\sqrt{(2\ell-1)(2\ell+3)}C^{\frac{3}{2}\frac{1}{2}2}_{\ell\ell j}\langle \ell, \ell_z; 2, \Delta\ell_z|\ell, \ell'_z\rangle\langle 3/2, s'_z; 2, \Delta\ell_z|1/2, s_z\rangle\ ,\end{aligned} \tag{39}$$

with the coefficients given in (37b).

The case $s' = 1/2$, $s = 3/2$ is completely analogous. From (3) we find

$$\begin{aligned}S(j, \ell, \ell, 1/2, 3/2) = &\frac{1}{2}\sqrt{\frac{3}{2}}C^{\frac{1}{2}\frac{3}{2}1}_{\ell\ell j}\langle \ell, \ell'_z|\widehat{\varepsilon}^k(\Delta\ell_z)L^k|\ell, \ell_z\rangle\widehat{\varepsilon}^h(\Delta\ell_z)^*\widehat{\chi}_A(s'_z)^*\widehat{\chi}^h_A(s_z)\\ &+\frac{1}{4}\sqrt{\frac{3}{2}}C^{\frac{1}{2}\frac{3}{2}2}_{\ell\ell j}\langle \ell, \ell'_z|\widehat{\varepsilon}^{k_1k_2}(\Delta\ell_z)L^{k_1}L^{k_2}|\ell, \ell_z\rangle\widehat{\varepsilon}^{h_1h_2}(\Delta\ell_z)^*\widehat{\chi}_A(s'_z)^*\sigma^{h_2}_{AB}\widehat{\chi}^{h_1}_B(s_z)\ ,\end{aligned} \tag{40}$$

with $C^{\frac{1}{2}\frac{3}{2}\Delta}_{\ell\ell j} = C^{\frac{3}{2}\frac{1}{2}\Delta}_{\ell\ell j}$ (see (37b)), and from (2)

$$\begin{aligned}S(j, \ell, \ell, 1/2, 3/2) = &\frac{1}{2}\sqrt{\frac{3}{2}}\sqrt{\ell(\ell+1)}C^{\frac{1}{2}\frac{3}{2}1}_{\ell\ell j}\langle \ell, \ell_z; 1, \Delta\ell_z|\ell, \ell'_z\rangle\langle 1/2, s'_z; 1, \Delta\ell_z|3/2, s_z\rangle\\ &+\frac{1}{8}\sqrt{\frac{5}{2}}\sqrt{\ell(\ell+1)}\sqrt{(2\ell-1)(2\ell+3)}C^{\frac{1}{2}\frac{3}{2}2}_{\ell\ell j}\langle \ell, \ell_z; 2, \Delta\ell_z|\ell, \ell'_z\rangle\langle 1/2, s'_z; 2, \Delta\ell_z|3/2, s_z\rangle\ .\end{aligned} \tag{41}$$

The expressions (37) and (40) will be used below to derive addition theorems involving one spin-3/2 and one spin-1/2 spherical harmonic.

$|\Delta\boldsymbol{\ell}| = \mathbf{1}$ We consider the case $s' = 3/2$, $s = 1/2$ first. In this case $S(j, \ell', \ell, 3/2, 1/2)$ can be non-vanishing only if $\ell' = \ell + 1$ and $j = \ell \pm 1/2$, or if $\ell' = \ell - 1$ and $j = \ell \pm 1/2$. With the spin matrix elements in (36), from (3) we obtain

$$\begin{aligned} S(j,\ell_1,\ell,3/2,1/2) = {} & \frac{1}{2}\sqrt{\frac{3}{2}} C^{\frac{3}{2}\frac{1}{2}1}_{\ell_1\ell j} \langle \ell_1 \ell'_z | \widehat{\varepsilon}^k(\Delta\ell_z) \widehat{r}^k | \ell, \ell_z \rangle \widehat{\varepsilon}^h(\Delta\ell_z)^* \widehat{\chi}^h_A(s'_z)^* \widehat{\chi}_A(s_z) \\ & + \frac{1}{4}\sqrt{\frac{3}{2}} C^{\frac{3}{2}\frac{1}{2}2}_{\ell_1\ell j} \langle \ell_1 \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) \widehat{r}^{k_1} L^{k_2} | \ell, \ell_z \rangle \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\chi}^{h_1}_A(s'_z)^* \sigma^{h_2}_{AB} \widehat{\chi}_B(s_z) \ , \end{aligned} \tag{42a}$$

with the coefficients

$$\begin{aligned} C^{\frac{3}{2}\frac{1}{2}1}_{\ell_1\ell j} &= -\sqrt{2}\Delta s \Delta\ell \sqrt{\frac{2j+1}{2\langle\ell\rangle+1}} \sqrt{\frac{2j - \langle\ell\rangle + \Delta s\Delta\ell + \frac{1}{2}}{2\langle\ell\rangle - \Delta s\Delta\ell + 1}} \sqrt{1 - 2\Delta s \Delta\ell (j - \langle\ell\rangle)} \ , \\ C^{\frac{3}{2}\frac{1}{2}2}_{\ell_1\ell j} &= -\frac{8}{\sqrt{3}} \sqrt{\frac{\frac{3}{2} + \Delta s\Delta\ell(j - \langle\ell\rangle)}{2\langle\ell\rangle - \Delta s\Delta\ell + 1}} \frac{1 + 2\Delta s\Delta\ell(j - \langle\ell\rangle)}{\sqrt{j + \frac{1}{2} + \Delta s\Delta\ell}} \ , \end{aligned} \tag{42b}$$

where in this case $\Delta s = 1$. On the first line of (42a) we can replace $\widehat{\varepsilon}^k(\Delta\ell_z)\widehat{\varepsilon}^h(\Delta\ell_z)^*$ with δ^{kh}, and on the second line,

$$\begin{aligned} \langle \ell' \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) \widehat{r}^{k_1} L^{k_2} | \ell, \ell_z \rangle \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\chi}^{h_1}_A(s'_z)^* \sigma^{h_2}_{AB} \widehat{\chi}_B(s_z) = \\ = \frac{1}{2} \langle \ell' \ell'_z | \widehat{r}^{\{k_1} L^{k_2\}_0} | \ell, \ell_z \rangle \widehat{\chi}^{h_1}_A(s'_z)^* \sigma^{h_2}_{AB} \widehat{\chi}_B(s_z) \ . \end{aligned} \tag{43}$$

Similarly, from (2) we get

$$\begin{aligned} S(j,\ell_1,\ell,3/2,1/2) = {} & -\frac{1}{2}\sqrt{\frac{3}{2}} \Delta\ell \sqrt{\frac{2\langle\ell\rangle+1}{2\ell_1+1}} C^{\frac{3}{2}\frac{1}{2}1}_{\ell_1\ell j} \langle \ell, \ell_z; 1, \Delta\ell_z | \ell_1, \ell'_z \rangle \langle 3/2, s'_z; 1, \Delta\ell_z | 1/2, s_z \rangle \\ & - \frac{\sqrt{30}}{32} \Delta\ell \sqrt{\frac{2\langle\ell\rangle+1}{2\ell_1+1}} \sqrt{(2\langle\ell\rangle - 1)(2\langle\ell\rangle + 3)} C^{\frac{3}{2}\frac{1}{2}2}_{\ell_1\ell j} \langle \ell, \ell_z; 2, \Delta\ell_z | \ell_1, \ell'_z \rangle \langle 3/2, s'_z; 2, \Delta\ell_z | 1/2, s_z \rangle \ , \end{aligned} \tag{44}$$

with the coefficients (42b).

In the case $s' = 1/2$, $s = 3/2$, $S(j, \ell', \ell, 1/2, 3/2)$ can be non-vanishing only if $\ell' = \ell + 1$ and $j = \ell + 3/2$, $\ell + 1/2$, or if $\ell' = \ell - 1$ and $j = \ell - 1/2$, $\ell - 3/2$. The treatment of this case is completely analogous to the previous one. In terms of matrix elements, from (3) we obtain

$$\begin{aligned} S(j,\ell_1,\ell,1/2,3/2) = {} & \frac{1}{2}\sqrt{\frac{3}{2}} C^{\frac{1}{2}\frac{3}{2}1}_{\ell_1\ell j} \langle \ell_1 \ell'_z | \widehat{\varepsilon}^k(\Delta\ell_z) \widehat{r}^k | \ell, \ell_z \rangle \widehat{\varepsilon}^h(\Delta\ell_z)^* \widehat{\chi}_A(s'_z)^* \widehat{\chi}^h_A(s_z) \\ & + \frac{1}{4}\sqrt{\frac{3}{2}} C^{\frac{1}{2}\frac{3}{2}2}_{\ell_1\ell j} \langle \ell_1 \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) \widehat{r}^{k_1} L^{k_2} | \ell, \ell_z \rangle \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\chi}_A(s'_z)^* \sigma^{h_1}_{AB} \widehat{\chi}^{h_2}_B(s_z) \ , \end{aligned} \tag{45}$$

with the coefficients $C^{\frac{1}{2}\frac{3}{2}\Delta}_{\ell_1\ell j}$ given by (42b) with $\Delta s = -1$. From (2) we get, analogously,

$$\begin{aligned} S(j,\ell_1,\ell,1/2,3/2) = {} & \frac{\sqrt{3}}{4} \Delta\ell \sqrt{\frac{2\langle\ell\rangle+1}{2\ell_1+1}} C^{\frac{1}{2}\frac{3}{2}1}_{\ell_1\ell j} \langle \ell, \ell_z; 1, \Delta\ell_z | \ell_1, \ell'_z \rangle \langle 1/2, s'_z; 1, \Delta\ell_z | 3/2, s_z \rangle \\ & + \frac{\sqrt{15}}{32} \Delta\ell \sqrt{\frac{2\langle\ell\rangle+1}{2\ell_1+1}} \sqrt{(2\langle\ell\rangle - 1)(2\langle\ell\rangle + 3)} C^{\frac{1}{2}\frac{3}{2}2}_{\ell_1\ell j} \langle \ell, \ell_z; 2, \Delta\ell_z | \ell_1, \ell'_z \rangle \langle 1/2, s'_z; 2, \Delta\ell_z | 3/2, s_z \rangle \ . \end{aligned} \tag{46}$$

The equalities (42) and (45) will be used for the derivation of an addition theorem involving one spin-3/2 and one spin-1/2 spherical harmonic, with orbital angular momentum differing by one unit.

$|\Delta\boldsymbol{\ell}| = \mathbf{2}$ As in the previous paragraph, we consider the case $s' = 3/2$, $s = 1/2$ first. In this case $S(j, \ell', \ell, 3/2, 1/2)$ can be non-vanishing only if $\ell' = \ell + 2$ and $j = \ell + 1/2$, or if $\ell' = \ell - 2$ and $j = \ell - 1/2$. Using the spin matrix elements (36) and the range of j, (3) reduces to

$$S(j, \ell_2, \ell, 3/2, 1/2) = \frac{1}{4}\sqrt{\frac{3}{2}} C^{\frac{3}{2}\frac{1}{2}2}_{\ell_2 \ell j} \langle \ell_2 \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) \widehat{r}^{k_1} \widehat{r}^{k_2} | \ell, \ell_z \rangle \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\chi}^{h_1}_A(s'_z)^* \sigma^{h_2}_{AB} \widehat{\chi}_B(s_z) \ , \tag{47a}$$

$$C^{\frac{3}{2}\frac{1}{2}2}_{\ell_2 \ell j} = \frac{2}{\sqrt{3}} \Delta s \Delta\ell \sqrt{2j+1} \sqrt{\frac{2\langle\ell\rangle + 1}{\langle\ell\rangle(\langle\ell\rangle + 1)}} \ , \tag{47b}$$

where $\Delta s = 1$. The matrix elements in (47a) can be written without standard tensors as,

$$\langle \ell_2 \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) \widehat{r}^{k_1} \widehat{r}^{k_2} | \ell, \ell_z \rangle \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\chi}^{h_1}_A(s'_z)^* \sigma^{h_2}_{AB} \widehat{\chi}_B(s_z) = \langle \ell_2 \ell'_z | \widehat{r}^{k_1} \widehat{r}^{k_2} | \ell, \ell_z \rangle \widehat{\chi}^{k_1}_A(s'_z)^* \sigma^{k_2}_{AB} \widehat{\chi}_B(s_z). \tag{48}$$

Similarly, in this case (2) takes the simplified form

$$\begin{aligned} S(j, \ell_2, \ell, 3/2, 1/2) = -\frac{\sqrt{30}}{8} \sqrt{2\ell+1} \sqrt{\frac{\langle\ell\rangle(\langle\ell\rangle+1)}{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)}} C^{\frac{3}{2}\frac{1}{2}2}_{\ell_2 \ell j} \\ \times \langle \ell, \ell_z; 2, \Delta\ell_z | \ell_2, \ell'_z \rangle \langle 3/2, s'_z; 2, \Delta\ell_z | 1/2, s_z \rangle \ . \end{aligned} \tag{49}$$

In the case $s' = 1/2$, $s = 3/2$, $S(j, \ell', \ell, 1/2, 3/2)$ can be non-vanishing only if $\ell' = \ell + 2$ and $j = \ell' - 1/2$, or if $\ell' = \ell - 2$ and $j = \ell' + 1/2$. In this case (3) reduces to

$$S(j, \ell_2, \ell, 1/2, 3/2) = \frac{1}{4}\sqrt{\frac{3}{2}} C^{\frac{1}{2}\frac{3}{2}2}_{\ell_2 \ell j} \langle \ell_2 \ell'_z | \widehat{\varepsilon}^{k_1 k_2}(\Delta\ell_z) \widehat{r}^{k_1} \widehat{r}^{k_2} | \ell, \ell_z \rangle \widehat{\varepsilon}^{h_1 h_2}(\Delta\ell_z)^* \widehat{\chi}_A(s'_z)^* \sigma^{h_2}_{AB} \widehat{\chi}^{h_1}_B(s_z) \ , \tag{50}$$

with $C^{\frac{1}{2}\frac{3}{2}2}_{\ell_2 \ell j}$ given by the same expression (47b) as $C^{\frac{3}{2}\frac{1}{2}2}_{\ell_2 \ell j}$, with $\Delta s = -1$. The equality (2) takes the simplified form

$$\begin{aligned} S(j, \ell_2, \ell, 1/2, 3/2) = \frac{\sqrt{15}}{8} \sqrt{2\ell+1} \sqrt{\frac{\langle\ell\rangle(\langle\ell\rangle+1)}{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)}} C^{\frac{1}{2}\frac{3}{2}2}_{\ell_2 \ell j} \\ \times \langle \ell, \ell_z; 2, \Delta\ell_z | \ell_2, \ell'_z \rangle \langle 1/2, s'_z; 2, \Delta\ell_z | 3/2, s_z \rangle \ , \end{aligned} \tag{51}$$

3 Angular momentum projection operators

We define the angular momentum projection operators as

$$\mathcal{P}_\ell = \sum_{m=-\ell}^{\ell} |\ell, m\rangle\langle\ell, m| \ , \qquad 0 \leq \ell < \infty \ . \tag{52}$$

It is immediate that $\mathcal{P}_\ell$ are a sequence of orthogonal projectors, which commute with $\vec{L}$ and resolve the identity,

$$\mathcal{P}_{\ell'} \mathcal{P}_\ell = \mathcal{P}_\ell \delta_{\ell'\ell} \ , \qquad \sum_{\ell=0}^{\infty} \mathcal{P}_\ell = 1 \ , \qquad [\vec{L}, \mathcal{P}_\ell] = 0 \ . \tag{53}$$

The matrix elements of $\mathcal{P}_\ell$ in configuration space follow from their definition (52) and the addition theorem for spherical harmonics

$$\mathcal{P}_\ell |\widehat{r}\,\rangle = \sum_{m=-\ell}^{\ell} Y_{\ell m}(\widehat{r})^* |\ell, m\rangle \ , \qquad \langle \widehat{r}\,' | \mathcal{P}_\ell | \widehat{r}\,\rangle = \frac{2\ell+1}{4\pi} P_\ell(\widehat{r} \cdot \widehat{r}\,') \ . \tag{54}$$

From (52)–(54) the matrix elements of $\mathcal{P}_\ell$ with other orbital operators can be computed. Those matrix elements, together with the results on products of CG coefficients of the previous section, are the basic building blocks needed in our derivation of addition theorems for spin spherical harmonics. As will be discussed in II, however, the results for $\mathcal{P}_\ell$ matrix elements constitute by themselves addition theorems for spherical harmonics.

3.1 Matrix elements with tensor powers of $\vec{L}$

The matrix elements of $\mathcal{P}_\ell$ with tensor powers of $\vec{L}$ will be used in the derivation of addition theorems for spin spherical harmonics. For the irreducible components, from which the full tensor matrix elements can be reconstructed, we have

$$\begin{aligned}\langle\widehat{r}\,'|L^{\{h_1}\dots L^{h_p\}_0}\mathcal{P}_\ell|\widehat{r}\rangle &= \frac{2\ell+1}{4\pi}i^p\,(\widehat{r}\wedge\nabla)^{\{h_p}\dots(\widehat{r}\wedge\nabla)^{h_1\}_0}\,P_\ell(x)\\ &= i\,(\widehat{r}\wedge\nabla)^{\{h_p}\,\langle\widehat{r}\,'|L^{h_1}\dots L^{h_{p-1}\}_0}\mathcal{P}_\ell|\widehat{r}\rangle\\ &= \frac{i}{(p-1)!}\,(\widehat{r}\wedge\nabla)^{\{h_p}\,\langle\widehat{r}\,'|L^{\{h_1}\dots L^{h_{p-1}\}_0\}_0}\mathcal{P}_\ell|\widehat{r}\rangle\ .\end{aligned}\tag{55}$$

Thus, we can obtain these irreducible matrix elements recursively, with the result

$$\langle\widehat{r}\,'|L^{\{h_1}\dots L^{h_p\}_0}\mathcal{P}_\ell|\widehat{r}\rangle = \frac{2\ell+1}{4\pi}i^p\sum_{q=0}^{[p/2]}C_{p,q}Z^{\{h_1\dots h_q;h_{q+1}\dots h_{2q}}v^{h_{2q+1}}\dots v^{h_p\}_0}P_\ell^{(p-q)}(x),\tag{56a}$$

with $P_\ell^{(k)}$ the k^{th} derivative of P_ℓ and,

$$\begin{gathered}x=\widehat{r}\cdot\widehat{r}\,'\ ,\quad \vec{v}=\widehat{r}\wedge\widehat{r}\,'\ ,\quad C_{p,q}=(2q-1)!!\binom{p}{2q}\ ,\quad C_{p,0}=1\ ,\\ Z^{i_1\dots i_{n_1};j_1\dots j_{n_2}}=\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_{n_1}}\widehat{r}\,'^{j_1}\dots\widehat{r}\,'^{j_{n_2}}\ .\end{gathered}\tag{56b}$$

For notational simplicity in what follows we adopt the following useful conventions. In $Z^{i_1\dots i_{n_1};j_1\dots j_{n_2}}$ one or both sets of indices may be empty,

$$Z^{\ ;j_1\dots j_{n_2}}=\widehat{r}\,'^{j_1}\dots\widehat{r}\,'^{j_{n_2}}\ ,\quad Z^{i_1\dots i_{n_2};\ }=\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_{n_1}}\ ,\quad Z^{\ ;\ }=1\ .\tag{56c}$$

In the term $q=0$ in (56a) the index sets are empty, $Z^{h_1\dots h_0;h_1\dots h_0}\equiv Z^{\ ;\ }$. Similarly, in the term $q=[p/2]$, when p is even, we set $v^{h_{2q+1}}\dots v^{h_p}=v^{h_{p+1}}\dots v^{h_p}\equiv 1$.

The cases $n=1$, 2, 3 will be needed for the derivation of addition theorems for spherical harmonics of spin 1/2, 1 and 3/2 so we quote them here explicitly,

$$\langle\widehat{r}\,'|\vec{L}\mathcal{P}_\ell|\widehat{r}\rangle = i\frac{2\ell+1}{4\pi}\vec{v}P_\ell'(\widehat{r}\cdot\widehat{r}\,')\ ,\tag{57a}$$

$$\langle\widehat{r}\,'|L^{\{i}L^{j\}_0}\mathcal{P}_\ell|\widehat{r}\rangle = -\frac{2\ell+1}{4\pi}\left(v^{\{i}v^{j\}_0}P_\ell''(\widehat{r}\cdot\widehat{r}\,')+\widehat{r}^{\,\{i}\widehat{r}\,'^{j\}_0}P_\ell'(\widehat{r}\cdot\widehat{r}\,')\right)\ ,\tag{57b}$$

$$\langle\widehat{r}\,'|L^{\{i}L^jL^{k\}_0}\mathcal{P}_\ell|\widehat{r}\rangle = -i\frac{2\ell+1}{4\pi}\left(v^{\{i}v^jv^{k\}_0}P_\ell'''(\widehat{r}\cdot\widehat{r}\,')+3\widehat{r}\,'^{\{i}\widehat{r}^{\,j}v^{k\}_0}P_\ell''(\widehat{r}\cdot\widehat{r}\,')\right)\ .\tag{57c}$$

3.2 Matrix elements with tensor powers of $\widehat{r}$

The matrix element $\langle\widehat{p}\,'|\mathcal{P}_{\ell'}\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_n}\mathcal{P}_\ell|\widehat{p}\rangle$ is an irreducible tensor if $\ell'=\ell\pm n$, otherwise reducible. The irreducible matrix elements will be used in the derivation of addition theorems for spin spherical harmonics. They satisfy the recursion relation,

$$\begin{aligned}&\langle\widehat{p}\,'|\mathcal{P}_{\ell_{n+1}}\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_{n+1}}\mathcal{P}_\ell|\widehat{p}\rangle =\\ &\qquad-\frac{1}{2\ell_n+1}\left[(\ell_{n+1}-\ell_n)\nabla_{p'}^{i_{n+1}}-\frac{1}{2}(\ell_{n+1}+\ell_n+1)\widehat{p}\,'^k\right]\langle\widehat{p}\,'|\mathcal{P}_{\ell_n}\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_n}\mathcal{P}_\ell|\widehat{p}\rangle\ ,\end{aligned}\tag{58}$$

with

$$\langle\widehat{p}\,'|\mathcal{P}_{\ell_1}\widehat{r}\,\mathcal{P}_\ell|\widehat{p}\rangle = \frac{\ell_1-\ell}{4\pi}\left(\widehat{p}\,'P_{\ell_1}'(x)-\widehat{p}\,P_\ell'(x)\right)\ .\tag{59}$$

In these equations we used the notation $\ell_k = \ell \pm k$, see appendix A. Eqs. (58), (59) are established in appendix D. The recursion relation (58) can be solved with the initial condition (59) to give

$$\begin{aligned}
\langle \widehat{p}\,'|\mathcal{P}_{\ell_n}\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_n}\mathcal{P}_\ell|\widehat{p}\rangle &= A_n \sum_{\substack{k_1,k_2=0\\ k_1+k_2=n}}^{n} \frac{(-1)^{k_2}}{k_1!k_2!} Z^{\{i_1\dots i_{k_2};i_{k_2+1}\dots i_n\}_0} P^{(n)}_{\ell_{k_1}}(x)\ , \\
A_n &= \frac{(\ell_1-\ell)^n}{4\pi}\frac{2\ell+1}{\prod_{k=0}^{n-1}(2\ell_k+1)} = \frac{1}{4\pi}\left(\frac{\ell_n-\ell}{n}\right)^n (2\ell+1)\frac{(\ell_n+\ell-n-\frac{\ell_n-\ell}{n})!!}{(\ell_n+\ell+n-\frac{\ell_n-\ell}{n})!!}\ ,
\end{aligned} \tag{60}$$

with $Z^{i_1\dots i_{n_1};j_1\dots j_{n_2}}$ defined in (56b), (56c). As before, we adopt the convention that in the term with $k_1 = n$, $k_2 = 0$ in (60) the first index set in Z is empty, $Z^{i_1\dots i_0;i_1\dots i_n} \equiv Z^{\ ;i_1\dots i_n}$ and, in the term with $k_1 = 0$, $k_2 = n$, the second index set is empty $Z^{i_1\dots i_n;i_{n+1}\dots i_n} \equiv Z^{i_1\dots i_n;\ }$.

The particular cases $n = 1$, 2, 3 will be needed for the derivation of addition theorems for spherical harmonics of spin 1/2, 1 and 3/2, so we give them here explicitly,

$$\langle \widehat{p}\,'|\mathcal{P}_{\ell_1}\widehat{r}^{\,i}\mathcal{P}_\ell|\widehat{p}\rangle = \frac{\ell_1-\ell}{4\pi}\left(-\widehat{p}^{\,i}P'_\ell(x) + \widehat{p}\,'^i P'_{\ell_1}(x)\right), \tag{61a}$$

$$\langle \widehat{p}\,'|\mathcal{P}_{\ell_2}\widehat{r}^{\,i_1}\widehat{r}^{\,i_2}\mathcal{P}_\ell|\widehat{p}\rangle = \frac{1}{4\pi}\frac{1}{2\ell_1+1}\left(\frac{1}{2}\widehat{p}^{\,\{i_1}\widehat{p}^{\,i_2\}_0}P''_\ell(x) - \widehat{p}^{\,\{i_1}\widehat{p}\,'^{i_2\}_0}P''_{\ell_1}(x) + \frac{1}{2}\widehat{p}\,'^{\{i_1}\widehat{p}\,'^{i_2\}_0}P''_{\ell_2}(x)\right), \tag{61b}$$

$$\begin{aligned}
\langle \widehat{p}\,'|\mathcal{P}_{\ell_3}\widehat{r}^{\,i_1}\widehat{r}^{\,i_2}\widehat{r}^{\,i_3}\mathcal{P}_\ell|\widehat{p}\rangle = \frac{\ell_1-\ell}{4\pi}\frac{1}{(2\ell_2+1)(2\ell_1+1)}\Big(&-\frac{1}{6}\widehat{p}^{\,\{i_1}\widehat{p}^{\,i_2}\widehat{p}^{\,i_3\}_0}P'''_\ell(x) + \frac{1}{2}\widehat{p}^{\,\{i_1}\widehat{p}^{\,i_2}\widehat{p}\,'^{i_3\}_0}P'''_{\ell_1}(x) \\
&-\frac{1}{2}\widehat{p}^{\,\{i_1}\widehat{p}\,'^{i_2}\widehat{p}\,'^{i_3\}_0}P'''_{\ell_2}(x) + \frac{1}{6}\widehat{p}\,'^{\{i_1}\widehat{p}\,'^{i_2}\widehat{p}\,'^{i_3\}_0}P'''_{\ell_3}(x)\Big).
\end{aligned} \tag{61c}$$

3.3 Mixed matrix elements

We consider now matrix elements of $\mathcal{P}_\ell$ with tensor products of both $\vec{L}$ and $\widehat{r}$. The matrix elements of the irreducible tensor operator $\widehat{r}^{\,\{i_1}\dots\widehat{r}^{\,i_n}L^{h_1}\dots L^{h_p\}_0}$ are of the form

$$\langle \widehat{p}\,'|\mathcal{P}_{\ell_n}\widehat{r}^{\,\{i_1}\dots\widehat{r}^{\,i_n}L^{h_1}\dots L^{h_t\}_0}\mathcal{P}_\ell|\widehat{p}\rangle = i^t(\widehat{p}\wedge\nabla_p)^{\{h_t}\dots(\widehat{p}\wedge\nabla_p)^{h_1}\langle \widehat{p}\,'|\mathcal{P}_{\ell_n}\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_n\}_0}\mathcal{P}_\ell|\widehat{p}\rangle\ . \tag{62}$$

From this equation and (60), for $t = 1$ we obtain

$$\langle \widehat{p}\,'|\mathcal{P}_{\ell_n}\widehat{r}^{\,\{i_1}\dots\widehat{r}^{\,i_n}L^{h\}_0}\mathcal{P}_\ell|\widehat{p}\rangle = iA_n\sum_{\substack{k_1,k_2=0\\ k_1+k_2=n}}^{n}(-1)^{k_2}\binom{n}{k_1}Z^{\{i_1\dots i_{k_2};i_{k_2+1}\dots i_n}v^{h\}_0}P^{(n+1)}_{\ell_{k_1}}(x)\ . \tag{63}$$

For $s > 1$, equation (62) leads to the recursion relation

$$\begin{aligned}
&\langle \widehat{p}\,'|\mathcal{P}_{\ell_n}\widehat{r}^{\,\{i_1}\dots\widehat{r}^{\,i_n}L^{h_1}\dots L^{h_t\}_0}\mathcal{P}_\ell|\widehat{p}\rangle = \\
&\qquad = \frac{i}{(n+t-1)!}(\widehat{p}\wedge\nabla_p)^{\{h_t}\langle \widehat{p}\,'|\mathcal{P}_{\ell_n}\widehat{r}^{\,\{i_1}\dots\widehat{r}^{\,i_n}L^{h_1}\dots L^{h_{t-1}\}_0\}_0}\mathcal{P}_\ell|\widehat{p}\rangle\ ,
\end{aligned} \tag{64}$$

which can be solved with the initial condition (63) to yield

$$\begin{aligned}
&\langle \widehat{p}\,'|\mathcal{P}_{\ell_n}\widehat{r}^{\,\{i_1}\dots\widehat{r}^{\,i_n}L^{h_1}\dots L^{h_t\}_0}\mathcal{P}_\ell|\widehat{p}\rangle = \\
&\quad = i^tA_n\sum_{\substack{k_1,k_2=0\\ k_1+k_2=n}}^{n}(-1)^{k_2}\binom{n}{k_1}\sum_{q=0}^{[t/2]}C_{t,q}Z^{\{i_1\dots i_{k_2}h_1\dots h_q;i_{k_2+1}\dots i_nh_{q+1}\dots h_{2q}}v^{h_{2q+1}}\dots v^{h_t\}_0}P^{(n+t-q)}_{\ell_{k_1}}(x)\ ,
\end{aligned} \tag{65}$$

with $[\]$ in the inner summation denoting integer part, A_n as defined in (60), and $C_{s,q}$, x, $\vec{v}$, and $Z^{i_1\dots i_{n_1};j_1\dots j_{n_2}}$ as defined in (56b), with $\widehat{p}$ instead of $\widehat{r}$. As before, for notational simplicity, we have not separated from the sum the terms with $(k_1,k_2) = (n,0)$, $(0,n)$, those with $q = 0$ and, for even s, those with $q = [s/2]$. In those cases we apply the same conventions as explained after eqs. (56) and (60).

Particular cases of importance for the derivation of addition theorems for spherical harmonics of spin 1/2, 1 and 3/2, are

$$\langle\widehat{p}'|\mathcal{P}_{\ell_1}\widehat{r}^{\{i}L^{k\}}\mathcal{P}_\ell|\widehat{p}\rangle = i\frac{\ell_1-\ell}{4\pi}\left(\widehat{p}'^{\{i}v^{k\}}P''_{\ell_1}(x) - \widehat{p}^{\{i}v^{k\}}P''_\ell(x)\right), \tag{66a}$$

$$\langle\widehat{p}'|\mathcal{P}_{\ell_2}\widehat{r}^{\{i}\widehat{r}^jL^{k\}_0}\mathcal{P}_\ell|\widehat{p}\rangle = \frac{i}{4\pi}\frac{1}{2\ell_1+1}\left(\widehat{p}'^{\{i}\widehat{p}'^jv^{k\}_0}P'''_{\ell_2}(x) - 2\widehat{p}'^{\{i}\widehat{p}^jv^{k\}_0}P'''_{\ell_1}(x) + \widehat{p}^{\{i}\widehat{p}^jv^{k\}_0}P'''_\ell(x)\right), \tag{66b}$$

$$\begin{aligned}\langle\widehat{p}'|\mathcal{P}_{\ell_1}\widehat{r}^{\{i}L^{j_1}L^{j_2\}_0}\mathcal{P}_\ell|\widehat{p}\rangle = -\frac{\ell_1-\ell}{4\pi}\Big(&\widehat{p}'^{\{i}v^{j_1}v^{j_2\}_0}P'''_{\ell_1}(x) - \widehat{p}^{\{i}v^{j_1}v^{j_2\}_0}P'''_\ell(x) + \widehat{p}'^{\{i}\widehat{p}'^{j_1}\widehat{p}^{j_2\}_0}P''_{\ell_1}(x)\\ &-\widehat{p}'^{\{i}\widehat{p}^{j_1}\widehat{p}^{j_2\}_0}P''_\ell(x)\Big).\end{aligned} \tag{66c}$$

4 Final remarks

We have presented in the foregoing sections preliminary results needed for the systematic derivation of addition theorems for spin spherical harmonics in II. In sect. 2 we obtained the factorization of orbital and spin degrees of freedom in products of CG coefficients of the form (1) in general form, and discussed the particular cases with $0 \le s', s \le 3/2$. In those cases the coefficients $C^{s's\Delta}_{\ell'\ell j}$, given in full generality in (3), were reduced to much smaller forms, and the tensor and spinor structures of each term in the expansion given explicitly.

In sect. 3 the matrix elements of the angular-momentum projector operator with the irreducible components of arbitrary tensor products of $\widehat{r}$ and $\vec{L}$ are given in general form, for all values of their parameters. Those matrix elements will be used in II to obtain general expressions for bilocal spherical harmonics, and to derive addition theorems for spin spherical harmonics. Their applicability is even wider, however, when they are appropriately combined to obtain matrix elements of reducible tensor operators. Some examples of those applications will also be considered in II.

As a side remark we point out that an unexpected byproduct of the results of sect. 2 is an improvement in computational efficiency, at least in the specific case of infinite-precision computation [13] in which the results are given as a rational number times the square root of a rational number. Consider, for example, the computation of both sides of (19), at fixed ℓ and j, for all possible values of $-\ell \le \ell'_z, \ell_z \le \ell$ and $-3/2 \le s'_z, s_z \le 3/2$. If the l.h.s. of (19) is computed as written on the second line of (1), we find that the ratio of CPU times $\tau_{\text{r.h.s.}}/\tau_{\text{l.h.s.}}$ begins at ~ 2 at $\ell = 1$, monotonically decreasing with ℓ to reach 1 at $\ell \sim 20$, ~ 0.5 at $\ell \sim 100$, and ~ 0.25 at $\ell \sim 200$. We remark that those numbers are subject to statistical fluctuations.

References

[1] A. R. Edmonds, "Angular Momentum in Quantum Mechanics," Princeton Univ. Press, Princeton, 1996.

[2] E. P. Wigner, "Group Theory and its Application to the Quantum Mechanics of Atomic Spectra," Academic Press, New York, 1959.

[3] B. R. Judd, "Angular Momentum Theory for Diatomic Molecules," Academic Press, New York, 1975.

[4] L. C. Biendenharn, J. D. Louck, "Angular Momentum in Quantum Physics," Encyclopedia of Mathemathics and its Applications Vol. 8, Cambridge Univ. Press, New York, 1985.

[5] L. C. Biendenharn, J. D. Louck, "The Racah-Wigner algebra in Quantum Theory," Encyclopedia of Mathemathics and its Applications Vol. 9, Cambridge Univ. Press, New York, 1985.

[6] D. A. Varshalovich, A. N. Moskalev, V. K. Khersonskii, "Quantum Theory of Angular Momentum," World Scientific, Singapore, 1988.

[7] M. Hamermesh, "Group Theory and its Application to Physical Problems," Dover, New York, 1989.

[8] A. Galindo, P. Pascual, "Quantum Mechanics," Vol. I, Springer–Verlag, New York, 1990.

[9] J. D. Jackson, "Classical Electrodynamics," John Wiley, New York, 1999.

[10] A. Bouzas, "Addition theorems for spin spherical harmonics. II Results," to be published.

[11] O. Nachtmann, "Elementary Particle Physics," Springer–Verlag, New York, 1990.

[12] Particle Data Group, C. Amsler et al., "Review of Particle Physics," Phys. Lett. B**667**, 2008, 1.

[13] S. Wolfram, "The Mathematica Book," Third Edition, Cambridge U. Press, New York, 1996.

[14] W. Rarita, J. Schwinger, Phys. Rev. **60**, 1941, 61.

A Notation and conventions

Throughout the paper we set $\hbar = 1$. We denote tensor product states of orbital and spin angular momentum by $|\ell, \ell_z; s, s_z\rangle = |\ell, \ell_z\rangle \otimes |s, s_z\rangle$, and states coupled to total angular momentum j, j_z by $|\ell, s, j, j_z\rangle$. We follow the notation of [7] for CG coefficients,

$$\langle \ell, \ell_z; s, s_z | j, j_z \rangle \equiv \langle \ell, \ell_z; s, s_z | \ell, s, j, j_z \rangle \ .$$

(Different notations are used, e.g., in [1, 6, 8, 12].) We adopt the usual convention that $\langle \ell, \ell_z; s, s_z | j, j_z \rangle = 0$ if $\ell_z + s_z \neq j_z$, or $j < |\ell - s|$, or $j > \ell + s$. For Wigner $6j$-symbols [1, 4–6] we use the definition and notation of [1, 6]. The most important property of $6j$-symbols for our purposes is the relation [1]

$$\begin{aligned} \langle j_1, m_1; j_2, m_2 | j', m_1 + m_2 \rangle \langle j', m_1 + m_2; j_3, m_3 | j, m \rangle = \sum_{j''=|j_2 - j_3|}^{j_2 + j_3} \sqrt{(2j'+1)(2j''+1)} (-1)^{j_1 + j_2 + j_3 + j} \\ \times \left\{ \begin{array}{ccc} j_1 & j_2 & j' \\ j_3 & j & j'' \end{array} \right\} \langle j_1, m_1; j'', m_2 + m_3 | j, m \rangle \langle j_2, m_2; j_3, m_3 | j'', m_2 + m_3 \rangle . \end{aligned} \tag{A.1}$$

This relation is satisfied by the implementation of CG coefficients and $6j$-symbols in the software system [13]. Our notation for the reduced matrix elements entering the WE theorem is explained in appendix B.1. For spherical harmonics $Y_{\ell m}$ and Legendre polynomials P_ℓ we use the standard definitions [1, 6, 8]. Furthermore, we adopt the convention, usual in numerical computations [13], $Y_{\ell m}(\widehat{r}\,) \equiv 0$ if $|m| > \ell$. When states with different angular momenta or spins are considered, we use the notations,

$$\langle \ell \rangle = \frac{\ell' + \ell}{2} \ , \quad \Delta\ell = \ell' - \ell \ , \quad \Delta\ell_z = \ell'_z - \ell_z \ , \tag{A.2}$$

and similarly $\langle s \rangle$, Δs, Δs_z. We also find useful the notation $\ell_k = \ell \pm k$ for orbital angular-momentum quantum numbers, with the convention that when several $\ell_{k_1}, \ldots, \ell_{k_p}$ appear in the same equation, then either the upper or the lower sign is chosen for all of them simultaneously. Thus, $\ell_{n+1} - \ell_n = \ell_{p+1} - \ell_p = \pm 1$ for any n and p.

We denote tensor indices by lower-case latin superindices, $A^{i_1 \ldots i_n}$, and spinor indices by upper-case latin subindices, χ_A. Since we consider only tensors and spinors of the su(2) algebra (as opposed to the sl(2) algebra), there is no need to raise spinor indices or lower tensor ones. We denote normal spinors and tensors by a caret, e.g., $\widehat{r} = \vec{r}/r$. Our choice of orthonormal bases for spinors and tensors is explained in appendix B.

Given a numeric or operator tensor $A^{i_1 \ldots i_n}$, we denote its associated totally symmetrized and antisymmetrized tensors as,

$$A^{\{i_1 \ldots i_n\}} = \sum_\sigma A^{i_{\sigma_1} \ldots i_{\sigma_n}} \ , \qquad A^{[i_1 \ldots i_n]} = \sum_\sigma \mathrm{sgn}(\sigma) A^{i_{\sigma_1} \ldots i_{\sigma_n}} \ , \tag{A.3}$$

where the sums extend over all permutations σ of $i_1, \ldots, i_n$. The same notations apply to tensor products, e.g., $\widehat{r}^{\,\{i} L^{j\}} = \widehat{r}^{\,i} L^j + \widehat{r}^{\,j} L^i$. We denote the traceless part of $A^{i_1 \ldots i_n}$ by $A^{(i_1 \ldots i_n)_0}$. The cases of importance in this paper are rank-2 and 3 tensors,

$$\begin{aligned} A^{(ij)_0} &= A^{ij} - \frac{1}{3} A^{kk} \delta^{ij} \ , \\ A^{(pqr)_0} &= A^{pqr} - \frac{2}{5} (A^{pjj} \delta^{qr} + A^{iqi} \delta^{pr} + A^{iir} \delta^{pq}) \\ &\quad + \frac{1}{10} (A^{qjj} \delta^{pr} + A^{rjj} \delta^{pq} + A^{ipi} \delta^{qr} + A^{iri} \delta^{pq} + A^{iip} \delta^{qr} + A^{iiq} \delta^{pr}) \ . \end{aligned} \tag{A.4}$$

The same notation applies to tensor products, such as $\widehat{r}^{(i}\widehat{r}^{j)_0} = \widehat{r}^i\widehat{r}^j - 1/3\delta^{ij}$. We denote by $A^{\{i_1\ldots i_n\}_0}$ the traceless part of $A^{\{i_1\ldots i_n\}}$. Thus, the irreducible component of $A^{i_1\ldots i_n}$ is $1/n!A^{\{i_1\ldots i_n\}_0}$. A more systematic approach to irreducible tensors is given in the following section.

B Standard tensor and spinor bases

In this appendix we introduce the standard bases for irreducible spinors and tensors of the su(2) algebra used as spin wavefunctions throughout the paper. We use spin wavefunctions of the Rarita-Schwinger type [14], which carry only vector indices varying from 1 to 3 and spinor indices from 1 to 2, and whose components are linearly related by symmetry and tracelessness and, for spinors, by transversality conditions. In fact, the spinors discussed in sect. B.2 are the non-relativistic version of Rarita-Schwinger spinors. The standard bases are defined so that they are orthonormal, have definite complex-conjugation properties and satisfy the Condon–Shortley phase conventions [1, 2, 4, 8].

Our starting point is the usual basis for Pauli spinors,

$$\widehat{\chi}_A(1/2) = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \ , \quad \widehat{\chi}_A(-1/2) = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \ . \tag{B.1}$$

The triplet component of basis-spinors tensor products yields a basis for the representation space of the adjoint representation,

$$\widehat{\chi}_{AB}(m) = \sum_{s'_z, s_z = -1/2}^{1/2} \langle \frac{1}{2}, s'_z; \frac{1}{2}, s_z | 1, m \rangle \widehat{\chi}(s'_z)_A \epsilon_{BC} \widehat{\chi}(s_z)_C \ , \quad m = 1, 2, 3 \ , \tag{B.2}$$

whose associated unit vectors are,

$$\widehat{\varepsilon}^k(1) = -\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \\ 0 \end{pmatrix} \ , \quad \widehat{\varepsilon}^k(0) = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \ , \quad \widehat{\varepsilon}^k(-1) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \\ 0 \end{pmatrix} \ . \tag{B.3}$$

These are the standard basis vectors, satisfying,

$$\widehat{\varepsilon}^k(s_z)^* = (-1)^{s_z} \widehat{\varepsilon}^k(-s_z) \ , \quad \widehat{\varepsilon}^k(s_z)^* \widehat{\varepsilon}^k(s'_z) = \delta_{s_z s'_z} \ , \quad \sum_{s_z=-1}^{1} \widehat{\varepsilon}^k(s_z) \widehat{\varepsilon}^{k'}(s_z)^* = \delta^{kk'} \ . \tag{B.4}$$

Furthermore, from the action of the spin operator on basis spinors we get

$$\widehat{\varepsilon}^k(m) \sigma^k_{AB} \widehat{\chi}_B(s_z) = \frac{2s_z - m}{\sqrt{1+m^2}} \widehat{\chi}_A((-1)^m s_z) \ , \quad m = -1, 0, 1 \ , s_z = \pm 1/2 \ , \tag{B.5}$$

a relation that will be needed in the discussion of spinors below.

B.1 Standard irreducible tensor bases

Starting with the standard vectors (B.3) we define the standard basis of irreducible rank-n tensors recursively as,

$$\widehat{\varepsilon}^{i_1\ldots i_n}(m) = \sum_{s_z=-1}^{1} \sum_{m'=-n+1}^{n-1} \langle n-1, m'; 1, s_z | n, m \rangle \widehat{\varepsilon}^{i_1\ldots i_{n-1}}(m') \widehat{\varepsilon}^{i_n}(s_z) \ , \quad -n \le m \le n \ . \tag{B.6}$$

This recursion can be solved to yield,

$$\widehat{\varepsilon}^{i_1\ldots i_n}(m) = \sum_{\substack{s_1,\ldots,s_n=-1 \\ s_1+\ldots+s_n=m}}^{1} f_n(s_1, \ldots, s_n) \widehat{\varepsilon}^{i_1}(s_1) \ldots \widehat{\varepsilon}^{i_n}(s_n) \ , \tag{B.7a}$$

with

$$f_n(s_1, \ldots, s_n) = \prod_{j=2}^{n} \langle 1, s_j; j-1, \sum_{i=1}^{j-1} s_i | j, \sum_{h=1}^{j} s_h \rangle = \left(\frac{2^n}{(2n)!} \frac{(n + \sum_{i=1}^n s_i)! (n - \sum_{j=1}^n s_j)!}{\prod_{h=1}^n (1+s_h)! (1-s_h)!} \right)^{\frac{1}{2}} \ . \tag{B.7b}$$

The second equality in (B.7b) follows from the first one by using the explicit expression for CG coefficients coupling angular momenta differing in one unit [7, 8]. From (B.7) it is clear that $\widehat{\varepsilon}^{i_1\ldots i_n}(m)$ is totally symmetric. Explicit evaluation shows that $\widehat{\varepsilon}^{i_1 i_2}(m)$, $-2 \le m \le 2$, are traceless and therefore it follows from (B.6) by induction that $\widehat{\varepsilon}^{i_1\ldots i_n}(m)$ are totally traceless for all $n \ge 2$, $-n \le m \le n$. Thus, (B.6) defines a set of $2n+1$ irreducible tensors of rank n. From (B.4) and (B.6) we obtain by induction the orthonormality and complex-conjugation relations

$$\widehat{\varepsilon}^{i_1\ldots i_n}(s_z)^* = (-1)^{s_z}\widehat{\varepsilon}^{i_1\ldots i_n}(-s_z) \ , \qquad \widehat{\varepsilon}^{i_1\ldots i_n}(s_z)^*\widehat{\varepsilon}^{i_1\ldots i_n}(s'_z) = \delta_{s_z s'_z} \ . \tag{B.8}$$

The completeness relation for the basis of standard tensors defines the orthogonal projector $X^{i_1\ldots i_n;j_1\ldots j_n}$ in the space of rank-n tensors onto the subspace of completely symmetric and traceless tensors

$$X^{i_1\ldots i_n;j_1\ldots j_n} = \sum_{s_z=-n}^{n} \widehat{\varepsilon}^{i_1\ldots i_n}(s_z)\widehat{\varepsilon}^{j_1\ldots j_n}(s_z)^* = \frac{1}{n!}\delta^{\{i_1 j_1}\ldots\delta^{i_n j_n\}'_0} \ . \tag{B.9a}$$

Here, the prime on the r.h.s. denotes symmetrization over the indices $i_1\ldots i_n$ and $j_1\ldots j_n$ separately, for example,

$$\begin{aligned}
X^{i;j} &= \delta^{ij} \ , \qquad X^{i_1 i_2;j_1 j_2} = \frac{1}{2}\left(\delta^{i_1 j_1}\delta^{i_2 j_2} + \delta^{i_1 j_2}\delta^{i_2 j_1} - \frac{2}{3}\delta^{i_1 i_2}\delta^{j_1 j_2}\right) \ , \\
X^{i_1 i_2 i_3;j_1 j_2 j_3} &= \frac{1}{6}\sum_{\sigma\{1,2,3\}}\Bigg[\delta^{i_1 j_{\sigma_1}}\delta^{i_2 j_{\sigma_2}}\delta^{i_3 j_{\sigma_3}} \\
&\quad - \delta^{i_1 i_2}\left(\frac{2}{5}\delta^{j_{\sigma_1} j_{\sigma_2}}\delta^{i_3 j_{\sigma_3}} - \frac{1}{10}\delta^{j_{\sigma_2} j_{\sigma_3}}\delta^{i_3 j_{\sigma_1}} - \frac{1}{10}\delta^{j_{\sigma_1} j_{\sigma_3}}\delta^{i_3 j_{\sigma_2}}\right) \\
&\quad - \delta^{i_1 i_3}\left(\frac{2}{5}\delta^{j_{\sigma_1} j_{\sigma_3}}\delta^{i_2 j_{\sigma_3}} - \frac{1}{10}\delta^{j_{\sigma_2} j_{\sigma_3}}\delta^{i_2 j_{\sigma_1}} - \frac{1}{10}\delta^{j_{\sigma_1} j_{\sigma_2}}\delta^{i_2 j_{\sigma_3}}\right) \\
&\quad - \delta^{i_2 i_3}\left(\frac{2}{5}\delta^{j_{\sigma_2} j_{\sigma_3}}\delta^{i_1 j_{\sigma_1}} - \frac{1}{10}\delta^{j_{\sigma_1} j_{\sigma_3}}\delta^{i_1 j_{\sigma_2}} - \frac{1}{10}\delta^{j_{\sigma_1} j_{\sigma_2}}\delta^{i_1 j_{\sigma_3}}\right)\Bigg] \ ,
\end{aligned} \tag{B.9b}$$

where in the last equality $\sigma\{1,2,3\}$ is the set of permutations of $\{1,2,3\}$. Using the recoupling identity (A.1) and (B.6) we can derive the important relation,

$$\widehat{\varepsilon}^{i_1\ldots i_n}(s_z) = \sum_{s_z=-q}^{q}\sum_{m'=-n+q}^{n-q}\langle n-q, m'; q, s_z|n, m\rangle\widehat{\varepsilon}^{i_1\ldots i_{n-q}}(m')\widehat{\varepsilon}^{i_{n-q+1}\ldots i_n}(s_z) \ , \quad 1 \le q \le n-1 \ , \tag{B.10}$$

showing that the maximal-rank coupling of two standard tensors is again a standard basis tensor. From (B.7) we obtain the equality

$$\widehat{\varepsilon}^{i_1\ldots i_n}(\pm n) = \widehat{\varepsilon}^{i_1}(\pm 1)\cdots\widehat{\varepsilon}^{i_n}(\pm 1) \ , \tag{B.11}$$

which is useful to evaluate reduced matrix elements. We adopt the convention

$$\widehat{\varepsilon}^{i_1\ldots i_n}(s_z) = 0 \quad \text{if} \quad |s_z| > n \ . \tag{B.12}$$

The matrix $\vec{S}_{(n)}$ of the spin operator $\vec{S}$ in the basis of spin-n states with n integer is given by

$$\begin{aligned}
\langle n, s'_z|S^j|n, s_z\rangle &= \widehat{\varepsilon}^{i_1\ldots i_n}(s'_z)^*(S^j_{(n)})^{i_1\ldots i_n;k_1\ldots k_n}\widehat{\varepsilon}^{k_1\ldots k_n}(s_z) \ , \\
(S^j_{(n)})^{i_1\ldots i_n;k_1\ldots k_n} &= \sum_{q=1}^{n} i\epsilon^{i_q j k_q}\prod_{\substack{p=1\\p\neq q}}^{n}\delta^{i_p k_n} \ .
\end{aligned} \tag{B.13}$$

If $A^{i_1\ldots i_n}$ is a rank-n real tensor, or a self-adjoint tensor operator, its irreducible component is

$$\frac{1}{n!}A^{\{i_1\ldots i_n\}_0} = \sum_{s_z=-n}^{n}\widehat{\varepsilon}^{i_1\ldots i_n}(s_z)\widehat{\varepsilon}^{j_1\ldots j_n}(s_z)^*A^{j_1\ldots j_n} = \sum_{s_z=-n}^{n}\widehat{\varepsilon}^{i_1\ldots i_n}(s_z)^*\widehat{\varepsilon}^{j_1\ldots j_n}(s_z)A^{j_1\ldots j_n} \ . \tag{B.14}$$

Furthermore, for self-adjoint tensor operators the WE theorem [8] holds,

$$\langle \ell', \ell'_z|\widehat{\varepsilon}^{i_1\ldots i_n}(s_z)A^{i_1\ldots i_n}|\ell, \ell_z\rangle = \langle \ell'||\widehat{\varepsilon}^{i_1\ldots i_n}A^{i_1\ldots i_n}||\ell\rangle\langle \ell, \ell_z; n, s_z|\ell', \ell'_z\rangle \ . \tag{B.15}$$

For a normal vector $\widehat{r}$, let $\mathcal{Y}_{nm}(\widehat{r}) = \widehat{\varepsilon}^{i_1\ldots i_n}(m)\widehat{r}^{\,i_1}\ldots\widehat{r}^{\,i_n}$. It follows from (B.6) by induction that

$$\vec{L}^2\mathcal{Y}_{nm}(\widehat{r}) = n(n+1)\mathcal{Y}_{nm}(\widehat{r}) \quad \text{and} \quad L^3\mathcal{Y}_{nm}(\widehat{r}) = m\mathcal{Y}_{nm}(\widehat{r}) \ . \tag{B.16}$$

We omit the proofs for brevity. We must then have $\mathcal{Y}_{nm}(\widehat{r}) \propto Y_{nm}(\widehat{r})$. Evaluation of both sides of the proportionality relation at $\widehat{r} = \widehat{z}$ yields

$$Y_{\ell m}(\widehat{r}) = \frac{1}{2\sqrt{\pi}}\sqrt{\frac{(2\ell+1)!!}{\ell!}}\widehat{\varepsilon}^{i_1\ldots i_\ell}(m)\widehat{r}^{\,i_1}\ldots\widehat{r}^{\,i_\ell} \ . \tag{B.17}$$

From this relation and the addition theorem for spherical harmonics we get

$$\begin{aligned}\widehat{r}^{\,(i_1}\ldots\widehat{r}^{\,i_\ell)_0}\widehat{r}^{\,\prime\,(i_1}\ldots\widehat{r}^{\,\prime\,i_\ell)_0} &= \sum_{m=-\ell}^{\ell}\widehat{\varepsilon}^{i_1\ldots i_\ell}(m)\widehat{\varepsilon}^{j_1\ldots j_\ell}(m)^*\widehat{r}^{\,i_1}\ldots\widehat{r}^{\,i_\ell}\widehat{r}^{\,\prime\,j_1}\ldots\widehat{r}^{\,\prime\,j_\ell} \\ &= \frac{\ell!}{(2\ell-1)!!}P_\ell(\widehat{r}\cdot\widehat{r}^{\,\prime}) \ .\end{aligned} \tag{B.18}$$

The relation (B.17) can be inverted, for each ℓ, by extending the basis (B.6) with appropriate reducible tensors to an orthogonal basis of the entire space of rank-n tensors.

B.2 Standard spinor bases

We define the standard spin-$(n+1/2)$ spinors with $n \geq 1$ as

$$\widehat{\chi}_A^{i_1\ldots i_n}(s_z) = \sum_{s_z'=-1/2}^{1/2}\sum_{m=-n}^{n}\langle n,m;1/2,s_z'|n+1/2,s_z\rangle\widehat{\varepsilon}^{i_1\ldots i_n}(m)\widehat{\chi}_A(s_z') \ , \qquad -n-1/2 \leq s_z \leq n+1/2 \ . \tag{B.19}$$

Using (B.6) and the recoupling identity (A.1), from (B.19) we find the equivalent expressions,

$$\widehat{\chi}_A^{i_1\ldots i_n}(s_z) = \sum_{s_z'=-q-1/2}^{q+1/2}\sum_{m=-n+q}^{n-q}\langle n-q,m;q+1/2,s_z'|n+1/2,s_z\rangle\widehat{\varepsilon}^{i_1\ldots i_{n-q}}(m)\widehat{\chi}_A^{i_{n-q+1}\ldots i_n}(s_z') \ , \tag{B.20}$$

with $1 \leq q \leq n-1$. From definition (B.19) we see that $\widehat{\chi}_A^{i_1\ldots i_n}$ is completely symmetric and traceless in its tensor indices. Thus, (B.20) remains valid after an arbitrary permutation of tensor indices in its r.h.s. From (B.19) with $n=1$ and (B.5) we find, for the spin-3/2 spinor, $\sigma_{AB}^i\widehat{\chi}_B^i(s_z) = 0$, $-3/2 \leq s_z \leq 3/2$. Thus, from (B.20) with $q=1$ we immediately obtain,

$$\sigma_{AB}^{i_k}\widehat{\chi}_B^{i_1\ldots i_k\ldots i_n}(s_z) = 0 \ , \qquad 1 \leq k \leq n \ , \quad -n-1/2 \leq s_z \leq n+1/2 \ . \tag{B.21}$$

From (B.19) and (B.21), the number of independent components in the spinor basis is $2n+2$ as it should. The orthornormality and complex-conjugation relations read

$$\widehat{\chi}_A^{i_1\ldots i_n}(s_z)^*\widehat{\chi}_A^{i_1\ldots i_n}(s_z') = \delta_{s_z s_z'} \ , \qquad i\sigma_{AB}^2\widehat{\chi}_B^{i_1\ldots i_n}(s_z) = (-1)^{1/2+s_z}\widehat{\chi}_A^{i_1\ldots i_n}(-s_z)^* \ . \tag{B.22}$$

The completeness relation for spin-s spinors, with $s = n+1/2$, $n \geq 1$, defines the orthogonal projector $X_{AB}^{i_1\ldots i_n;j_1\ldots j_n}$ onto the subspace of spin-$(n+1/2)$ spinors of $\mathbb{C}^{(2n+1)\times 2}$, [2]

$$\begin{aligned}X_{AB}^{i_1\ldots i_n;j_1\ldots j_n} &\equiv \sum_{s_z=-s}^{s}\widehat{\chi}_A^{i_1\ldots i_n}(s_z)\widehat{\chi}_B^{j_1\ldots j_n}(s_z)^* \\ &= X^{i_1\ldots i_n;k_1\ldots k_n}\left[\frac{n+1}{2n+1}\delta^{k_1h_1}\ldots\delta^{k_nh_n}\delta_{AB} + \frac{n}{2n+1}i\epsilon^{k_1rh_1}\sigma_{AB}^r\delta^{k_2h_2}\ldots\delta^{k_nh_n}\right]X^{i_1\ldots i_n;k_1\ldots k_n} \ ,\end{aligned} \tag{B.23}$$

[2] For notational simplicity we identify the space of complex rank-n completely symmetric and traceless tensors with $\mathbb{C}^{(2n+1)}$, which has the same dimension.

where $X^{i_1\ldots i_n;j_1\ldots j_n}$ is the projector defined in (B.9). The orthogonal complement of the spin-$(n+1/2)$ subspace in $\mathbb{C}^{(2n+1)\times 2}$ is $\ker(X)$, which can be parameterized as $\sigma^{\{i_1}_{AB}\widehat{\chi}^{i_2\ldots i_n\}_0}_B$ with $\widehat{\chi}^{i_1\ldots i_{(n-1)}}_A$ a spin-$(n-1/2)$ spinor. This is the spin-$(n-1/2)$ subspace of $\mathbb{C}^{(2n+1)\times 2}$.

The spin operator for $s=(n+1/2)$ spinors is given by the operator tensor product of the spin-n and spin-1/2 operators, its matrix being

$$(S^j_{(n+1/2)})^{i_1\ldots i_n;k_1\ldots k_n}_{AB} = (S^j_{(n)})^{i_1\ldots i_n;k_1\ldots k_n}\delta_{AB} + \frac{1}{2}\delta^{i_1k_1}\ldots\delta^{i_nk_n}\sigma^j_{AB}\ , \tag{B.24}$$

with $S^j_{(n)}$ defined in (B.13). From (B.21) and (B.24) we obtain the following useful identities,

$$i\epsilon^{i_1kh}\sigma^k_{AB}\widehat{\chi}^{hi_2\ldots i_n}_B(s_z) = \widehat{\chi}^{i_1\ldots i_n}_A(s_z)\ , \tag{B.25}$$

$$\sigma^k_{AB}\widehat{\chi}^{i_1\ldots i_n}_B(s_z) - \sigma^{i_1}_{AB}\widehat{\chi}^{ki_2\ldots i_n}_B(s_z) + i\epsilon^{ki_1h}\widehat{\chi}^{hi_2\ldots i_n}_A(s_z) = 0\ , \tag{B.26}$$

$$\widehat{\chi}^{i_1\ldots i_{n-1}h}_A(s'_z)^*\sigma^k_{AB}\widehat{\chi}^{j_1\ldots j_{n-1}h}_B(s_z) = \widehat{\chi}^{i_1\ldots i_{n-1}p}_A(s'_z)^* i\epsilon^{pkq}\widehat{\chi}^{j_1\ldots j_{n-1}q}_A(s_z)\ , \tag{B.27}$$

$$\begin{aligned}\widehat{\chi}^{i_1\ldots i_n}_A(s'_z)^*(S^k_{(n+1/2)})^{i_1\ldots i_n;j_1\ldots j_n}_{AB}\widehat{\chi}^{j_1\ldots j_n}_B(s_z) &= (n+1/2)\widehat{\chi}^{h_1\ldots h_n}_A(s'_z)^*\sigma^k_{AB}\widehat{\chi}^{h_1\ldots h_n}_B(s_z)\\ &= (n+1/2)\widehat{\chi}^{h_1\ldots h_{n-1}p}_A(s'_z)^* i\epsilon^{pkq}\widehat{\chi}^{h_1\ldots h_{n-1}q}_B(s_z)\ ,\end{aligned} \tag{B.28}$$

which remain valid after permutation of $i_1\ldots i_n$, etc.

C Reduced matrix elements

In this appendix we gather expressions for matrix elements of certain irreducible tensor operators needed in the foregoing. The notation used is the same as in (B.15). We begin from the known results [8]

$$\langle j'||\widehat{\varepsilon}^i J^i||j\rangle = \sqrt{j(j+1)}\delta_{jj'}\ , \tag{C.1}$$

$$\langle \ell'||Y_L||\ell\rangle = \frac{1}{\sqrt{4\pi}}\sqrt{\frac{(2\ell+1)(2L+1)}{2\ell'+1}}\langle \ell,0;L,0|\ell',0\rangle\ . \tag{C.2}$$

In (C.1) J^i can of course be any angular momentum operator: orbital L^i, spin S^i or total J^i. For tensor products of angular momentum operators we have, using (B.11) and (C.1),

$$\langle j'||\widehat{\varepsilon}^{i_1\ldots i_n}J^{i_1}\ldots J^{i_n}||j\rangle = \frac{n!}{\sqrt{2^n(2n)!}}\frac{1}{\sqrt{2j+1}}\sqrt{\frac{(2j+n+1)!}{(2j-n)!}}\,\delta_{jj'}\ . \tag{C.3}$$

For convenience, we quote here the explicit form of the CG coefficient appearing in (C.2). If t is odd the CG coefficient vanishes, and if $t\geq 0$ is even

$$\begin{aligned}\langle \ell,0;n+t,0|\ell_n,0\rangle = (-1)^{t/2}(\ell_1-\ell)^n&\sqrt{\frac{(2n+t-1)!!(t-1)!!}{(n+t/2)!(t/2)!}}\sqrt{2\ell_n+1}\sqrt{\frac{((\ell_n+\ell+n+t)/2)!}{((\ell_n+\ell-n-t)/2)!}}\\ &\times\sqrt{\frac{(\ell_n+\ell-n-t-1)!!}{(\ell_n+\ell+n+t+1)!!}}\ ,\qquad t\geq 0 \text{ even.}\end{aligned} \tag{C.4}$$

Next, we consider tensor products of the position versor $\widehat{r}=\vec{r}/r$. From (B.17) and (C.2)

$$\langle \ell'||\widehat{\varepsilon}^{i_1\ldots i_n}\widehat{r}^{i_1}\ldots\widehat{r}^{i_n}||\ell\rangle = \sqrt{\frac{n!}{(2n-1)!!}}\sqrt{\frac{2\ell+1}{2\ell'+1}}\langle \ell,0;n,0|\ell',0\rangle\ . \tag{C.5}$$

In the case $|\Delta\ell|=n$, which is of interest to us, a more explicit expression is given by,

$$\begin{aligned}\langle \ell_n||\widehat{\varepsilon}^{i_1\ldots i_n}\widehat{r}^{i_1}\ldots\widehat{r}^{i_n}||\ell\rangle = \frac{(-1)^{\frac{n-\Delta\ell}{2}}}{2^{n/2}}&\sqrt{\frac{2\ell+1}{2(\langle\ell\rangle+\frac{n}{2})+1}}\sqrt{\frac{\Gamma\left(\langle\ell\rangle-\frac{n}{2}+\frac{1}{2}\right)}{\Gamma\left(\langle\ell\rangle+\frac{n}{2}+\frac{1}{2}\right)}}\\ &\times\sqrt{\frac{\Gamma\left(\langle\ell\rangle+\frac{n}{2}+1\right)}{\Gamma\left(\langle\ell\rangle-\frac{n}{2}+1\right)}}\ .\end{aligned} \tag{C.6}$$

The cases $n = 1, 2, 3$ are of particular importance to us. Noticing that $(-1)^{\frac{|\Delta\ell|-\Delta\ell}{2}} = \left(\frac{\Delta\ell}{|\Delta\ell|}\right)^{|\Delta\ell|}$, we get

$$\langle \ell_1 || \widehat{\varepsilon}^i \widehat{r}^i || \ell \rangle = \frac{\Delta\ell}{\sqrt{2}} \sqrt{\frac{2\langle\ell\rangle+1}{2\ell_1+1}} \,, \tag{C.7a}$$

$$\langle \ell_2 || \widehat{\varepsilon}^{i_1 i_2} \widehat{r}^{i_1} \widehat{r}^{i_2} || \ell \rangle = \sqrt{2\ell+1} \sqrt{\frac{\langle\ell\rangle(\langle\ell\rangle+1)}{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)}} \,, \tag{C.7b}$$

$$\langle \ell_3 || \widehat{\varepsilon}^{i_1 i_2 i_3} \widehat{r}^{i_1} \widehat{r}^{i_2} \widehat{r}^{i_3} || \ell \rangle = \frac{\Delta\ell}{24\sqrt{2}} \sqrt{\frac{2\ell+1}{\langle\ell\rangle+2}} \sqrt{\frac{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)}{(\langle\ell\rangle-1)\langle\ell\rangle(\langle\ell\rangle+1)}} \,. \tag{C.7c}$$

Equations (C.6) and (C.7) remain valid under the replacements $\ell \to s$, $\widehat{r}^i \to T^i$, that give the reduced matrix elements $\langle s \pm n || \widehat{\varepsilon}^{i_1 \cdots i_n} T^{i_1} \dots T^{i_n} || s \rangle$ of tensor products of the spin transition operator for integer or half-integer s.

For tensor products of $\widehat{r}$ and $\vec{L}$, using (C.3), (C.6) and (B.11) we obtain,

$$\begin{aligned} \langle \ell + \Delta\ell || \widehat{\varepsilon}^{i_1 \cdots i_n} \widehat{r}^{i_1} \dots \widehat{r}^{i_{|\Delta\ell|}} L^{i_{|\Delta\ell|+1}} \dots L^{i_n} || \ell \rangle = \left(\frac{\Delta\ell}{|\Delta\ell|}\right)^{|\Delta\ell|} \frac{1}{2^{n/2} 2^{|\Delta\ell|}} \sqrt{\frac{(n+|\Delta\ell|)!(n-|\Delta\ell|)!}{(2n)!}} \\ \times \frac{\sqrt{2\ell+1}}{2\langle\ell\rangle+|\Delta\ell|+1} \sqrt{\frac{(2\langle\ell\rangle+n+1)!}{(2\langle\ell\rangle-n)!}} \frac{\Gamma\left(\langle\ell\rangle-|\Delta\ell|/2+1/2\right)}{\Gamma\left(\langle\ell\rangle+|\Delta\ell|/2+1/2\right)} \,. \end{aligned} \tag{C.8}$$

This equation reduces to (C.3) for $\Delta\ell = 0$, and to (C.6) for $|\Delta\ell| = n$. The particular cases $n = 2, 3$ of (C.8) are important in the foregoing,

$$\langle \ell_1 || \widehat{\varepsilon}^{ij} \widehat{r}^i L^j || \ell \rangle = \frac{\Delta\ell}{4} \frac{1}{\sqrt{2\ell_1+1}} \sqrt{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)} \,, \tag{C.9a}$$

$$\langle \ell_1 || \widehat{\varepsilon}^{ijk} \widehat{r}^i L^j L^k || \ell \rangle = \frac{\Delta\ell}{2\sqrt{30}} \frac{1}{\sqrt{2\ell_1+1}} \sqrt{(2\langle\ell\rangle-2)(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)(2\langle\ell\rangle+4)} \,, \tag{C.9b}$$

$$\langle \ell_2 || \widehat{\varepsilon}^{ijk} \widehat{r}^i \widehat{r}^j L^k || \ell \rangle = \frac{1}{\sqrt{3}} \sqrt{2\ell+1} \sqrt{\frac{(\langle\ell\rangle-1)\langle\ell\rangle(\langle\ell\rangle+1)(\langle\ell\rangle+2)}{(2\langle\ell\rangle-1)(2\langle\ell\rangle+1)(2\langle\ell\rangle+3)}} \,. \tag{C.9c}$$

The following mixed reduced matrix elements are also useful when dealing with tensor decompositions in irreducible components,

$$\langle \ell_1 || \widehat{\varepsilon}^i (\widehat{r} \wedge \vec{L})^i || \ell \rangle = -\frac{i}{2\sqrt{2}} \left((\ell_1 + \ell + 1) - 2\Delta\ell \right) \sqrt{\frac{\ell_1+\ell+1}{2\ell_1+1}} \,, \tag{C.10a}$$

$$\langle \ell_1 || \widehat{\varepsilon}^{ij} (\widehat{r} \wedge \vec{L})^i L^j || \ell \rangle = -\frac{i}{8} \left((\ell_1 + \ell + 1) - 2\Delta\ell \right) \sqrt{\frac{(\ell_1+\ell-1)(\ell_1+\ell+1)(\ell_1+\ell+3)}{2\ell_1+1}} \,, \tag{C.10b}$$

$$\langle \ell_2 || \widehat{\varepsilon}^{ij} \widehat{r}^i (\widehat{r} \wedge \vec{L})^j || \ell \rangle = -\frac{i}{2} \left((\ell_1 - \ell)(2\ell_1+1) - 3 \right) \sqrt{\frac{\ell_1(\ell_1+1)}{(2\ell_1+1)(2\ell_2+1)}} \,, \tag{C.10c}$$

With the replacements $\ell \to s$, $\Delta\ell \to \Delta s$, $\widehat{r}^i \to T^i$, $L^i \to S^i$, (C.8) and (C.9) give the reduced matrix elements $\langle s \pm n || \widehat{\varepsilon}^{i_1 \cdots i_n} T^{i_1} \dots T^{i_{|\Delta s|}} S^{i_{|\Delta s|+1}} \dots S^{i_n} || s \rangle$ for integer or half-integer s, and similarly for (C.10).

D The matrix elements $\langle \widehat{\boldsymbol{p}}' | \mathcal{P}_{\ell_n} \widehat{\boldsymbol{r}}^{\boldsymbol{i_1}} \dots \widehat{\boldsymbol{r}}^{\boldsymbol{i_n}} \mathcal{P}_{\boldsymbol{\ell}} | \widehat{\boldsymbol{p}} \rangle$

In this appendix we derive (58) and (59). First, we consider the matrix element,

$$\begin{aligned} \langle \widehat{p}' | \mathcal{P}_{\ell_1} \widehat{p}' \cdot \widehat{r}\, \mathcal{P}_\ell | \widehat{p} \rangle &= \int d^2q\, \langle \widehat{p}' | \mathcal{P}_{\ell_1} | \widehat{q} \rangle \widehat{p}' \cdot \widehat{q}\, \langle \widehat{q} | \mathcal{P}_\ell | \widehat{p} \rangle \\ &= \frac{2\ell_1+1}{4\pi} \frac{2\ell+1}{4\pi} \int d^2q\, P_{\ell_1}(\widehat{p}' \cdot \widehat{q}) \widehat{p}' \cdot \widehat{q}\, P_\ell(\widehat{p} \cdot \widehat{q}) \\ &= \frac{\ell_1+1}{2\ell_1+3} \langle \widehat{p}' | \mathcal{P}_{\ell_1+1} \mathcal{P}_\ell | \widehat{p} \rangle + \frac{\ell_1}{2\ell_1-1} \langle \widehat{p}' | \mathcal{P}_{\ell_1-1} \mathcal{P}_\ell | \widehat{p} \rangle \,, \end{aligned} \tag{D.1}$$

where the integrals extend over the unit sphere, and in the last equality we used the relation

$$P_{\ell_1}(x) = \frac{\ell_1+1}{2\ell_1+1}P_{\ell_1+1}(x) + \frac{\ell_1}{2\ell_1-1}P_{\ell_1-1}(x) \ .$$

Applying the projector property of $\mathcal{P}_\ell$ in the last line of (D.1) we obtain,

$$\langle\widehat{p}\,'|\mathcal{P}_{\ell_1}\widehat{p}\,'\cdot\widehat{r}\,\mathcal{P}_\ell|\widehat{p}\rangle = \frac{1}{2}\frac{1}{2\ell+1}(\ell_1+\ell+1)\langle\widehat{p}\,'|\mathcal{P}_\ell|\widehat{p}\rangle \ . \tag{D.2}$$

Similarly, using

$$\nabla_{p'}\widehat{p}\,'\cdot\widehat{q} = \widehat{p}\,'\wedge(\widehat{q}\wedge\widehat{p}\,') \ ,$$

we get,

$$\langle\widehat{p}\,'|\mathcal{P}_{\ell_1}\widehat{p}\,'\wedge(\widehat{r}\wedge\widehat{p}\,')\mathcal{P}_\ell|\widehat{p}\rangle = \nabla_{p'}\left(\frac{\ell_1-\ell}{4\pi}P_{\ell_1}(\widehat{p}\cdot\widehat{p}\,')\right) = -\frac{\ell_1-\ell}{4\pi}\widehat{p}\,'\wedge(\widehat{p}\wedge\widehat{p}\,')P'_\ell(\widehat{p}\cdot\widehat{p}\,') \ . \tag{D.3}$$

Summing (D.2) and (D.3) we obtain (59) with the help of the relation

$$xP'_{\ell_k}(x) - (n-k)P_{\ell_k}(x) + \frac{1}{2}(\ell_{n+1}-\ell_n)(\ell_{n+1}+\ell_n+1)P_{\ell_k} = P'_{\ell_{k+1}} \ , \quad 0\le k\le n \ . \tag{D.4}$$

The derivation of (58) runs along the same lines,

$$\begin{aligned}\langle\widehat{p}\,'|\mathcal{P}_{\ell_{n+1}}\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_n}\widehat{r}^{\,i_{n+1}}\mathcal{P}_\ell|\widehat{p}\rangle &= \int d^2q\,\langle\widehat{p}\,'|\mathcal{P}_{\ell_{n+1}}\widehat{r}^{\,i_{n+1}}\mathcal{P}_{\ell_n}|\widehat{q}\rangle\langle\widehat{q}\,|\mathcal{P}_{\ell_n}\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_n}\mathcal{P}_\ell|\widehat{p}\rangle \\ &= \int d^2q\,\langle\widehat{p}\,'|\mathcal{P}_{\ell_{n+1}}\widehat{p}\,'^{i_{n+1}}\widehat{p}\,'\cdot\widehat{r}\,\mathcal{P}_{\ell_n}|\widehat{q}\rangle\langle\widehat{q}\,|\mathcal{P}_{\ell_n}\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_n}\mathcal{P}_\ell|\widehat{p}\rangle \\ &+ \int d^2q\,\langle\widehat{p}\,'|\mathcal{P}_{\ell_{n+1}}(\widehat{r}^{\,i_{n+1}}-\widehat{p}\,'^{i_{n+1}}\widehat{p}\,'\cdot\widehat{r}\,)\mathcal{P}_{\ell_n}|\widehat{q}\rangle\langle\widehat{q}\,|\mathcal{P}_{\ell_n}\widehat{r}^{\,i_1}\dots\widehat{r}^{\,i_n}\mathcal{P}_\ell|\widehat{p}\rangle \ .\end{aligned} \tag{D.5}$$

Applying now (D.2) to the first integral and (D.3) to the second one in the last equality, we obtain (58).